

Symplectic extension graphs and Ribet's lemma

Aniruddha Sudarshan

ABSTRACT. Let p be an odd prime, and let $\rho: G \rightarrow \mathrm{GSp}_{2n}(\mathbb{Z}_p)$ be an irreducible symplectic residually multiplicity free representation. We define the symplectic extension graph $\Gamma_\rho^{\mathrm{GSp}}$ associated with ρ to be the directed graph whose vertices are the residual factors of ρ and a directed edge between two vertices corresponds to a nonsplit extension between the vertices realized by ρ . Using the Euclidean Bruhat–Tits building associated with $\mathrm{GSp}_{2n}(\mathbb{Q}_p)$, under some assumptions, we prove the existence of certain paths in $\Gamma_\rho^{\mathrm{GSp}}$ and show that $\Gamma_\rho^{\mathrm{GSp}}$ is a disjoint union of nonempty strongly connected subgraphs. Moreover, we provide examples to show that the result we obtain is optimal.

CONTENTS

1. Introduction	1
2. Symplectic modules and representations	6
3. Symplectic types and constructing idempotents	11
4. Buildings	18
5. Symplectic extension graph	30
References	36

1. Introduction

In his remarkable paper, Ribet proves the converse to Herbrand's theorem by using certain congruences between modular forms [Rib76]. A crucial step in his proof is the construction of a nonsplit extension of characters, usually called *Ribet's lemma*. Throughout this paper, we fix an odd prime p . Let V be a 2-dimensional $\mathbb{Q}_p$ -vector space, $\rho: G \rightarrow \mathrm{GL}(V)$ be an irreducible representation that admits a G -stable $\mathbb{Z}_p$ -lattice in V , and let ρ be residually reducible with factors χ_1 and χ_2 . Ribet shows the existence of G -stable lattices Λ_1 and Λ_2 of V such that $\Lambda_i/p\Lambda_i$ is a nonsplit extension of χ_i by χ_j for all $1 \leq i \neq j \leq 2$ [Rib76, Proposition 2.1].

The Galois representation associated with a cuspidal elliptic Hecke eigenform that is congruent to an Eisenstein series modulo p satisfies Ribet's hypothesis. Using the local

2020 *Mathematics Subject Classification.* Primary 20G25; Secondary 51E24, 20C11.

Key words and phrases. Ribet's lemma, Bruhat–Tits buildings, Galois representations.

properties (such as *ordinary* at p) of such a Galois representation, Ribet constructs nontrivial unramified cohomology classes. This method of using congruences between modular forms to construct nontrivial elements in certain Selmer groups is often called *Ribet's method*.

1.1. Result of Bellaïche. Bellaïche attributes to Serre a proof of [Rib76, Proposition 2.1] using the Bruhat–Tits tree associated with $\mathrm{GL}_2(\mathbb{Q}_p)$ [Bel09]. Generalizing this method, for all $n \geq 2$, Bellaïche obtains a GL_n -version of [Rib76, Proposition 2.1] using the Bruhat–Tits building associated with $\mathrm{GL}_n(\mathbb{Q}_p)$ [Bel03, Theorem 1].

Let V be a finite dimensional vector space over $\mathbb{Q}_p$, and let $\rho: G \rightarrow \mathrm{GL}(V)$ be a p -adic representation that admits a G -stable $\mathbb{Z}_p$ -lattice in V . For example, this occurs when G is compact and ρ is continuous [Bel09, Corollary 1.1]. For a G -stable lattice Λ , let $\bar{\rho}_\Lambda$ be the residual representation $\Lambda/p\Lambda$. Although the representation $\bar{\rho}_\Lambda$ depends on Λ , its semisimplification $\bar{\rho}_\Lambda^{\mathrm{ss}}$ is independent of the choice of Λ [Bel09, Proposition 1.2]. The *residual factors* of a representation $\rho: G \rightarrow \mathrm{GL}(V)$ that admits a G -stable lattice Λ are defined to be the irreducible Jordan–Hölder factors of $\bar{\rho}_\Lambda^{\mathrm{ss}}$. We assume the residual factors of $\bar{\rho}$ are absolutely irreducible. Let $\rho_1, \dots, \rho_d$ be the isomorphism classes of the residual factors of ρ .

DEFINITION 1.1. The *extension graph* $\Gamma_\rho^{\mathrm{GL}}$ is the directed graph on the set of vertices $\{\rho_1, \dots, \rho_d\}$, where an edge from ρ_i to ρ_j exists if there exists a G -stable lattice Λ in V such that $\bar{\rho}_\Lambda$ admits a nonsplit extension of ρ_i by ρ_j as a subquotient.

A directed graph is *strongly connected* if there is a directed path between any two vertices.

THEOREM 1.2. [Bel03, Theorem 1] *If ρ is irreducible, then $\Gamma_\rho^{\mathrm{GL}}$ is strongly connected.*

Under the assumption that each ρ_i occurs with multiplicity one in $\bar{\rho}^{\mathrm{ss}}$, Bellaïche–Graftieaux show that ρ is irreducible if and only if $\Gamma_\rho^{\mathrm{GL}}$ is strongly connected [BG06, Proposition 3.4.1]. Thus, Theorem 1.2 is optimal under the given hypothesis. When $d = 2$, Theorem 1.2 is a reformulation of [Rib76, Proposition 2.1]. Using Ribet's method, Theorem 1.2 has some arithmetic applications (for example, [BC04] and [BK21]).

1.2. Our results. We define and study a symplectic variant of Bellaïche's extension graph. Let V be a $\mathbb{Q}_p$ -vector space with a symplectic (see Definition 2.1) bilinear form ϕ .

DEFINITION 1.3. A $\mathbb{Z}_p$ -lattice Λ in V is a *symplectic lattice* if the bilinear form on $\Lambda/p\Lambda$ induced¹ by ϕ is symplectic. If Λ is symplectic and $\phi(\Lambda \times \Lambda) = \mathbb{Z}_p$, then we call Λ *primitive*.

Let $\mathrm{GL}(V, \phi)$ be the subgroup of $\mathrm{GL}(V)$ consisting of elements g such that,

$$\phi(gv, gw) = \lambda(g)\phi(v, w) \quad \text{for all } v, w \in V \text{ and for some } \lambda(g) \in \mathbb{Q}_p^\times.$$

A representation $\rho: G \rightarrow \mathrm{GL}(V)$ is *symplectic* if $\rho(G) \subseteq \mathrm{GL}(V, \phi)$.

Let $\rho: G \rightarrow \mathrm{GL}(V, \phi)$ be an irreducible symplectic representation that admits a G -stable primitive lattice. We note that ρ stabilizes a primitive lattice if and only if there exists a (symplectic) basis of V with respect to which $\rho(G) \subseteq \mathrm{GSp}_{2n}(\mathbb{Z}_p)$. We assume the residual

¹See the discussion before Definition 4.3

factors of $\bar{\rho}$ are absolutely irreducible. Let $\rho_1, \dots, \rho_d$ be the isomorphism classes of the residual factors of ρ .

DEFINITION 1.4. The *symplectic extension graph* Γ_ρ^{GSp} is the directed graph on the set of vertices $\{\rho_1, \dots, \rho_d\}$ such that an edge from ρ_i to ρ_j exists if there exists a G -stable symplectic lattice Λ in V such that $\bar{\rho}_\Lambda$ admits a nonsplit extension of ρ_i by ρ_j as a subquotient.

We show that Γ_ρ^{GSp} is a *skew symmetric graph* (see Definition 1.5). Under further assumptions on the residual factors of ρ , we show the existence of certain paths in Γ_ρ^{GSp} and show that Γ_ρ^{GSp} is a disjoint union of strongly connected graphs.

To state our results precisely, we introduce some notation. For all $g \in G$, we let $\lambda_\rho(g) := \lambda(\rho(g))$. Since ρ admits a G -stable primitive lattice, we see that $\lambda_\rho(G) \subseteq \mathbb{Z}_p^\times$. Let τ be the $\mathbb{Z}_p$ -linear involution of the group ring $\mathbb{Z}_p[G]$ defined on G by $\tau(g) = \lambda_\rho(g)g^{-1}$. For a matrix M , we let M^t be its transpose. For a representation $\psi: G \rightarrow \text{GL}_{2n}(\mathbb{Z}_p)$, we let $\psi^\perp := (\psi \circ \tau)^t$. Since $\rho(G) \subseteq \text{GL}(V, \phi)$, there exists a transposition σ on $I = \{1, \dots, d\}$ such that

$$\rho_{\sigma(i)} \simeq \rho_i^\perp \quad \text{for all } i \in I.$$

DEFINITION 1.5. [GK96] A directed graph (V, E) is *skew symmetric* if there is a transposition σ on V such that $(u, v) \in E$ if and only if $(\sigma(v), \sigma(u)) \in E$.

PROPOSITION 1.6. The directed graph Γ_ρ^{GSp} is a skew symmetric graph with respect to the transposition σ on $\{\rho_1, \dots, \rho_d\}$ defined by $\sigma(\rho_i) := \rho_{\sigma(i)}$ for all $i \in I$.

Throughout the paper, we usually assume

(rmf) ρ is residually multiplicity free.

For our main results, we also need an extra assumption:

(dist) There exists $g_0 \in G$ such that $\bar{\rho}(g_0)$ has distinct eigenvalues.

We now state the main theorem of the paper. All the results we state for p -adic representations hold true, with the same proof, when we replace $\mathbb{Z}_p$ by a complete discrete valuation ring with finite residue field of characteristic p .

THEOREM 1.7. Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space. Let $\rho: G \rightarrow \text{GL}(V, \phi)$ be an irreducible symplectic representation that admits a G -stable primitive lattice, with absolutely irreducible residual factors $\rho_1, \dots, \rho_d$. Let σ be the permutation on $\{1, \dots, d\}$ defined by $\rho_{\sigma(i)} \simeq \rho_i^\perp$. Assume that ρ satisfies (dist) and (rmf), and that $\sigma(i) \neq i$ for all $1 \leq i \leq d$. Then the directed graph Γ_ρ^{GSp} satisfies the following conditions.

- (a) For all $1 \leq i \leq d$, ρ_i and $\rho_{\sigma(i)}$ lie in the same strongly connected component.
- (b) It is a disjoint union of at most $d/2$ strongly connected graphs.

In particular, if Γ_ρ^{GSp} has $d/2$ strongly connected components, then Γ_ρ^{GSp} is the disjoint union of strongly connected graphs with vertices $\{\rho_i, \rho_{\sigma(i)}\}_{i < \sigma(i)}$.

When $d = 2$, Theorem 1.7 is a generalization of [Urb01, Proposition 1.1] (Urban works over GSp_4) for complete discrete valuation rings with finite residue field of odd characteristic.

Using [Urb01, Proposition 1.1], Urban produces nontrivial elements in a Selmer group of certain twists of ad^0 of a modular p -adic Galois representation [Urb01, Theorem 3.7].

We show that Theorem 1.7 is the optimal result one could achieve in this generality.

THEOREM 1.8. *Given a positive integer n , and an integer $r \in \{1, \dots, n\}$, there exists a group G_r and an irreducible symplectic representation $\rho: G_r \hookrightarrow \text{GSp}_{2n}(\mathbb{Z}_p)$ satisfying (dist) and (rmf) such that*

- (a) *the residual factors of ρ are characters,*
- (b) *Γ_ρ^{GSp} is a disjoint union of r strongly connected graphs.*

We believe (dist) should be omittable, and should be equivalent to (rmf) (this is the case when $n = 1$). On the other hand, the assumption $\sigma(i) \neq i$ for all $1 \leq i \leq d$ is necessary for Theorem 1.7 and the method of proof. To be precise, we produce a symplectic irreducible p -adic representation ρ that is residually reducible with three absolutely irreducible residual factors such that Γ_ρ^{GSp} is an empty graph but Γ_ρ^{GL} is a complete graph (see Proposition 5.14).

1.2.1. The work of Bellaïche–Graftieaux. In proving our main results, we prove symplectic versions of certain results in [BG06]. In this subsection, we give an overview of such results.

Let V be an n -dimensional $\mathbb{Q}_p$ -vector space and let $\rho: G \rightarrow \text{GL}(V)$ be an irreducible representation that admits a G -stable $\mathbb{Z}_p$ -lattice. Suppose that ρ satisfies (rmf) and $\rho_1, \dots, \rho_d$ are its absolutely irreducible residual factors. Let S_ρ^{GL} be the (finite) set homothety classes of G -stable lattices in the Bruhat–Tits building associated with $\text{GL}(V)$. Then there exists a basis $\{v_1, \dots, v_n\}$ of V such that every vertex in S_ρ^{GL} is of the form $[a_1, \dots, a_n] := [\langle p^{a_1}v_1, \dots, p^{a_n}v_n \rangle]$ for some integers $a_1, \dots, a_n$ [BG06, Proposition 3.1.1] (see Proposition 4.26 for the analogous symplectic version). Moreover, there exists a metric $\eta: \{1, \dots, n\}^2 \rightarrow \mathbb{Z}$ such that

$$S_\rho^{\text{GL}} = \{[a_1, \dots, a_n] \mid a_i - a_j \leq \eta(i, j)\}.$$

In fact, S_ρ^{GL} lives in a sub-apartment of dimension $d - 1$: there exists a partition $\{1, \dots, n\} = \bigsqcup_{j=1}^d B_j$, such that $\eta = 0$ on $B_j \times B_j$ for all $1 \leq j \leq d$. This implies that η is constant on $B_j \times B_k$ for all $1 \leq j, k \leq d$. Thus, η defining S_ρ^{GL} defines a metric graph $\mathcal{G}(S_\rho^{\text{GL}})$ on $\{1, \dots, d\}$. Conversely, given a metric graph $\mathcal{G}$ on $\{1, \dots, d\}$, there exists a group G and an irreducible representation $\rho: G \rightarrow \text{GL}_n(\mathbb{Z}_p)$ such that $\mathcal{G}(S_\rho^{\text{GL}}) = \mathcal{G}$ [BG06, Lemme 3.2.1]. The proof of Theorem 1.8 is analogous to the proof of [BG06, Lemme 3.2.1].

1.3. Method of proof. Let V be a finite dimensional $\mathbb{Q}_p$ -vector space with a symplectic form ϕ . Let $\rho: G \rightarrow \text{GL}(V, \phi)$ be an irreducible symplectic representation that admits a G -stable primitive lattice and satisfies (rmf). Let $\rho_1, \dots, \rho_d$ be the absolutely irreducible residual factors of ρ . Let $I = \{1, \dots, d\}$, $\sigma: I \rightarrow I$ be defined by $\rho_{\sigma(i)} \simeq \rho_i^\perp$, and let $I_0 := \{i \in I \mid \sigma(i) = i\}$.

We prove Theorem 1.7 by using the Bruhat–Tits building $\Delta(V, \phi)$ associated with $\text{GL}(V, \phi)$. The building $\Delta(V, \phi)$ is a simplicial complex whose vertices are either homothety classes $[\Lambda_0]$ of primitive lattices, or correspond to isotropic subspaces of $\Lambda_0/p\Lambda_0$. Let $\mathcal{S}_\rho^{\text{GSp}}$ be the simplicial subcomplex of $\Delta(V, \phi)$ formed by G -stable vertices in $\Delta(V, \phi)$. Since ρ is irreducible, $\mathcal{S}_\rho^{\text{GSp}}$ is compact. The vertices on the boundary of $\mathcal{S}_\rho^{\text{GSp}}$ correspond to indecomposable residual representations. Under the assumption $I_0 = \emptyset$, we show the existence of certain

symplectic vertices on the boundary of $\mathcal{S}_\rho^{\text{GSp}}$. These are precisely the vertices that realize certain edges in the symplectic extension graph associated with ρ .

The assumption $I_0 = \emptyset$ is equivalent to the existence of neighbouring symplectic vertices in $\mathcal{S}_\rho^{\text{GSp}}$ (see [Proposition 4.27](#)). We define the Jordan–Hölder colouring on the edges in $\mathcal{S}_\rho^{\text{GSp}}$ as in [\[Bel03\]](#). Assuming $I_0 = \emptyset$ and [\(dist\)](#), we can move between symplectic vertices in $\mathcal{S}_\rho^{\text{GSp}}$ to show the existence of G -stable symplectic vertices (on the boundary) with no edges of a specific colouring (see [Theorem 4.34](#)). This is equivalent to the existence of edges in Γ_ρ^{GSp} (see [Proposition 4.33](#)).

The proof of [Theorem 1.8](#) builds on the examples constructed when $d \in \{2, 4\}$. To prove [Theorem 1.8](#), we use a construction inspired by [\[BG06, Lemme 3.2.1\]](#). Given a positive integer $r \leq n$, we define a metric $\eta^{(r)}$ on I such that $G_r := \{(a_{ij}) \in \text{GSp}_{2n}(\mathbb{Z}_p) \mid a_{ij} \in p^{\eta^{(r)}(i,j)}\mathbb{Z}_p\}$ with the inclusion $\rho: G_r \hookrightarrow \text{GSp}_{2n}(\mathbb{Z}_p)$ satisfies the conclusions of [Theorem 1.8](#). A similar construction is used to produce an example of a p -adic representation whose symplectic extension graph is empty (see [Proposition 5.14](#)).

1.4. Motivation and future direction. Ribet's lemma was generalized to prove certain cases of the Iwasawa main conjectures [\[MW84, Wil90, Urb01, SU14\]](#). More recently, Dasgupta–Kakde and collaborators proved the Brumer–Stark conjecture by generalizing Ribet's lemma for two-dimensional representations valued in complete reduced local rings [\[DK23, DKS23a, DKS23b\]](#).

Bellaïche–Chenevier generalize Ribet's lemma to local henselian rings by use of *pseudorepresentations* and *generalized matrix algebras* (GMAs) [\[BC09, §1\]](#). Given an irreducible pseudocharacter that is residually multiplicity free, they provide a systematic way to construct nonsplit extensions between residual representations [\[BC09, Theorem 1.5.5\]](#). As an arithmetic application, they construct more than one linearly independent element in certain Selmer groups to prove cases of the Bloch–Kato conjecture [\[BC09, Chapter 5\]](#).

Recently, Moakher–Quast have developed an analogous theory of symplectic *determinants* (in the sense of [\[Che14\]](#)) and symplectic GMAs [\[MQ26\]](#). Building on this new theory, one hopes for a symplectic analogue, if it exists, of [\[BC09, Theorem 1.5.5\]](#) and its possible arithmetic applications. As a first step in this pursuit, we study extensions of representations realized by symplectic lattices.

1.5. Outline of the paper. In [Section 2](#), we fix notation and provide background on symplectic modules and representations. In [Section 3](#), we recall a few results and definitions from [\[BC09, §1.8\]](#) in our setting and prove [Proposition 1.6](#). The key result of this section is, given a symplectic residually multiplicity-free p -adic representation ρ , under some assumptions, we show the existence of a “special” symplectic basis with respect to which the diagonal matrices lie in the image of ρ .

In [Section 4](#), we introduce Bruhat–Tits buildings associated with general linear groups and symplectic groups over $\mathbb{Q}_p$. A building is an abstract simplicial complex that is a union of subcomplexes called apartments. By their description, an apartment corresponds to a basis of the underlying vector space and is easier to work with. For example, moving between vertices in an apartment corresponds to conjugation of lattices by certain diagonal matrices. For a symplectic irreducible p -adic representation ρ , we study the geometry of $\mathcal{S}_\rho^{\text{GSp}}$, the simplicial

complex formed by G -stable lattices in the symplectic building. We show that $\mathcal{S}_\rho^{\text{GSp}}$ lies in a symplectic apartment corresponding to a special symplectic basis as in [Section 3](#). This gives us all the data of vertices and their neighbours in $\mathcal{S}_\rho^{\text{GSp}}$ with respect to the symplectic basis, and is used to prove the existence of certain boundary vertices of $\mathcal{S}_\rho^{\text{GSp}}$.

In [Section 5](#), using the main results from [Section 4](#), we prove our main results [Theorem 1.7](#) and [Theorem 1.8](#). We end with an example of an irreducible symplectic representation whose extension graph is an empty graph (see [Proposition 5.14](#)).

Acknowledgement. I would like to thank my advisor Jackie Lang for continuous support and encouragement throughout writing this article. I would also like to thank Mohamed Moakher for many useful conversations and comments on an earlier draft, Matthew Stover for conversations on buildings, and Carl Wang-Erickson for helpful suggestions.

2. Symplectic modules and representations

In this section, we give a brief introduction to symplectic modules and representations.

2.1. Symplectic modules. We recall a few facts about finite free symplectic modules over a commutative ring A with $2 \in A^\times$. Given an A -basis $\{e_1, \dots, e_n\}$ of V , the matrix representing ϕ on V is the $(n \times n)$ matrix whose (i, j) -th entry is $\phi(e_i, e_j)$.

DEFINITION 2.1. A *symplectic pairing* on a finite free A -module V is a bilinear pairing $\phi: V \times V \rightarrow A$ satisfying

- (1) (alternating) $\phi(v, v) = 0$ for all $v \in V$;
- (2) (perfect) the map $V \mapsto \text{Hom}_A(V, A)$ mapping v to the A -linear map $w \mapsto \phi(v, w)$ is an isomorphism of A -modules.

REMARKS.

- (a) A bilinear pairing ϕ is a symplectic pairing on V if and only if, for any given basis of V , the matrix $Q \in M_n(A)$ representing ϕ is invertible and skew symmetric; that is, $Q \in \text{GL}_n(A)$ and $Q^t = -Q$. Using this, one can check that a symplectic pairing on a finite free A -module V exists if and only if V has even rank.
- (b) [Definition 2.1\(2\)](#) is stronger than just ϕ being nondegenerate. To be precise, if ϕ is a symplectic pairing on a finite free module V over a ring A , then for every nonzero $w \in V$, there exists $v \in V$ such that $\phi(v, w) = 1$.
- (c) When A is a field, [Definition 2.1\(2\)](#) is equivalent to ϕ being nondegenerate.

By the notion of a symplectic pairing, we make the following definitions as in [\[LS10, §1\]](#).

DEFINITIONS 2.2. A tuple (V, ϕ) is called a *symplectic module* over A of rank $2n$ if V is a finitely generated free module over A with rank $2n$ and ϕ is a symplectic bilinear pairing on V . A submodule W of V is said to be

- (1) *isotropic* if $\phi|_{W \times W} = 0$;
- (2) *symplectic* if $\phi|_{W \times W}$ is nondegenerate.

We see that a $\mathbb{Z}_p$ -lattice Λ in a symplectic $\mathbb{Q}_p$ -vector space (V, ϕ) is primitive (that is, $\phi(\Lambda \times \Lambda) = \mathbb{Z}_p$ and $\phi|_{\Lambda \times \Lambda} \bmod p$ is symplectic on $\Lambda/p\Lambda$) if and only if Λ is a symplectic

$\mathbb{Z}_p$ -module with respect to $\phi|_{\Lambda \times \Lambda}$. By definition, if Λ is a symplectic $\mathbb{Z}_p$ -module, then it is primitive. We will see later (see [Corollary 4.11](#)) that a primitive $\mathbb{Z}_p$ -lattice is a symplectic $\mathbb{Z}_p$ -module with respect to $\phi|_{\Lambda \times \Lambda}$ by [Proposition 2.5](#) below.

DEFINITION 2.3. For a subset W of V , we write W^0 for the submodule of V consisting of elements $w^0 \in V$ such that $\phi(w^0, w) = 0$ for all $w \in W$.

We observe that W is isotropic (respectively, symplectic) if and only if $W \subseteq W^0$ (respectively, $W \cap W^0 = 0$).

DEFINITION 2.4. An isotropic subspace W of V is *Lagrangian* if it satisfies $W = W^0$. In other words, a Lagrangian subspace is a maximal isotropic subspace of V .

PROPOSITION 2.5. Let V be a free module of rank $2n$ over a ring A with $2 \in A^\times$. Let ϕ be an alternating bilinear pairing on V . Then ϕ is a symplectic pairing if and only if, given a nonzero element $w_1 \in V$, there is a basis of V containing w_1 with respect to which the matrix representing ϕ is

$$J_{2n} := \begin{pmatrix} & -I_n \\ I_n & \end{pmatrix},$$

where I_n is the identity matrix in $\text{GL}_n(A)$.

PROOF. The *if* part is immediate. We now prove the *only if* part. By definition, there exists $v_1 \in V$ such that $\phi(v_1, w_1) = 1$. If $n = 1$, then V is the span of $\{w_1, v_1\}$ and the result follows. Let W be the span of $\{w_1, v_1\}$. Then it is easy to see that $V = W \oplus W^0$ and W^0 is a symplectic submodule of V of dimension $2n - 2$. The proof follows by inducting on n . $\square$

When the dimension n is implicit, we write J and I instead of J_{2n} and I_n , respectively.

DEFINITION 2.6. Let (V, ϕ) be a symplectic module over a ring A with $2 \in A^\times$. A basis of V is a *symplectic basis* if the matrix representing ϕ is J .

LEMMA 2.7. Let (V, ϕ) be a symplectic module of rank $2n$ over a ring A with $2 \in A^\times$. Given a symplectic basis B of V , and an integer $1 \leq r \leq n$, there exists a basis of V , given by permuting elements of B , with respect to which the matrix representing ϕ is

$$\begin{pmatrix} & -I_r \\ & J_{2(n-r)} \\ I_r & \end{pmatrix}.$$

PROOF. Let $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ be a symplectic basis of V as constructed in [Proposition 2.5](#). The required basis is $\{w_1, \dots, w_n, v_{r+1}, \dots, v_n, v_1, \dots, v_r\}$. $\square$

2.1.1. Properties of symplectic subspaces. In this section, we recall a few basic facts on the symplectic subspaces of a symplectic vector space (V, ϕ) over a field K with $\text{Char}(K) \neq 2$.

PROPOSITION 2.8. If W is a symplectic subspace of (V, ϕ) , then

$$V = W \oplus W^0.$$

PROOF. We choose a symplectic basis $\{w_1, \dots, w_r, v_1, \dots, v_r\}$ of W by [Proposition 2.5](#). For $v \in V$, we have $\tilde{v} \in W^0$ and $v - \tilde{v} \in W$. This shows that $W + W^0 = V$. The result now follows since W is symplectic and $W \cap W^0 = 0$. $\square$

COROLLARY 2.9. *A subspace W of (V, ϕ) is symplectic if and only if W^0 is symplectic. Moreover, if W is symplectic, then $(W^0)^0 = W$.*

PROOF. If W is symplectic, then it follows that W^0 is symplectic by [Proposition 2.8](#). For the converse, if W is not symplectic, then there is nonzero $w \in W \cap W^0$. If W^0 were symplectic, then there exists $v \in W^0$ such that $\phi(w, v) = 1$. Since $w \in W$ and $v \in W^0$, we have a contradiction. This proves that if W^0 is symplectic, then so is W . If W and W^0 are symplectic, then by looking at dimensions, it follows that $W = (W^0)^0$. $\square$

COROLLARY 2.10. *Let (V, ϕ) be a symplectic vector space of dimension $2n$. Let W be a symplectic subspace of V of dimension $2r$. Any symplectic basis of W can be extended to a basis of V with respect to which the matrix corresponding to ϕ is*

$$\begin{pmatrix} J_{2r} & \\ & J_{2(n-r)} \end{pmatrix}.$$

PROOF. We see that $V = W \oplus W^0$ and both W and W^0 are symplectic subspaces of V by [Proposition 2.8](#) and [Corollary 2.9](#). There exists a symplectic basis B of W and a symplectic basis B^0 of W^0 by [Proposition 2.5](#). The desired basis is $B \sqcup B^0$. $\square$

2.1.2. *Properties of isotropic subspaces.* Let (V, ϕ) be a symplectic vector space of dimension $2n$ over a field K with $\text{Char}(K) \neq 2$.

LEMMA 2.11. *Let W be an isotropic subspace of V . There is a basis of W that extends to a symplectic basis of V . Moreover, the dimension of W is n if and only if W is a Lagrangian subspace of V .*

PROOF. The proof of the first part follows by induction on $\dim_K W$ for which the base case is [Proposition 2.5](#). If W is an isotropic subspace of dimension r , then W is Lagrangian inside a $2r$ -dimensional symplectic subspace of V . This proves the second part. $\square$

The following result generalizes [Proposition 2.5](#) and is crucial. An isotropic flag in a symplectic vector space corresponds to a simplex in the corresponding symplectic building. We use the following [Proposition 2.12](#) in [Section 4](#) to show that every simplex lies in an apartment (see [Proposition 4.12](#)).

PROPOSITION 2.12. *Let (V, ϕ) be a symplectic vector space over a field K with $\text{Char}(K) \neq 2$, and let*

$$0 = V_0 \subset V_1 \subset \dots \subset V_m \subset V$$

be a flag in V such that V_m is isotropic. There is a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of V such that $\{w_1, \dots, w_{i_k}\}$ is a basis of V_k for $1 \leq k \leq m$, where $i_k = \dim V_k$.

PROOF. We induct on $m \geq 1$. The base case is [Lemma 2.11](#). Using the inductive hypothesis on the flag $0 \subset V_1 \subset \dots \subset V_{m-1} \subset V$, we get a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$

of V such that $\{w_1, \dots, w_{i_k}\}$ is a basis for V_k for $1 \leq k < m$. For $w \in V_m$, consider the element in V_m defined by $\tilde{w} = w + \sum_{j=1}^{i_{m-1}} \phi(w, v_j) w_j$. By our choice, $\phi(\tilde{w}, v_j) = 0$ for $1 \leq j \leq i_{m-1}$. Let $\tilde{V}_m = \{\tilde{w} \mid w \in V_m\}$. Since $\tilde{V}_m$ is an isotropic subspace of the symplectic space $\{w_1, \dots, w_{i_{m-1}}, v_1, \dots, v_{i_{m-1}}\}^0$, using the induction hypothesis, we obtain a symplectic basis of $\{w_1, \dots, w_{i_{m-1}}, v_1, \dots, v_{i_{m-1}}\}^0$ extending a basis $\{w_{i_{m-1}+1}, \dots, w_{i_m}\}$ of $\tilde{V}_m$. By combining the two bases suitably, the result follows. $\square$

2.2. Symplectic representations. In this section, we study the properties of symplectic representations. Using this, we give a proof of [Proposition 1.6](#) (see [Section 5](#)).

DEFINITION 2.13. Let A be a commutative ring with $2 \in A^\times$. A *morphism* between symplectic modules (V, ϕ) and (V', ϕ') over A is an A -module morphism $\sigma: V \rightarrow V'$ such that, for all $v, w \in V$,

$$\phi'(\sigma(v), \sigma(w)) = \lambda(\sigma)\phi(v, w) \quad \text{for some } \lambda(\sigma) \in A.$$

A morphism $\sigma: (V, \phi) \rightarrow (V', \phi')$ is an *isomorphism* if there exists a two-sided inverse morphism $\sigma^{-1}: (V', \phi') \rightarrow (V, \phi)$.

Note that symplectic morphisms are closed under composition. Moreover, for symplectic morphisms $\sigma: V \rightarrow V'$ and $\tau: V' \rightarrow V''$, we see that $\lambda(\tau \circ \sigma) = \lambda(\tau)\lambda(\sigma)$.

DEFINITION 2.14. For a symplectic module (V, ϕ) over A , let $\text{End}_A(V, \phi)$ be the monoid (with respect to composition) of endomorphisms of (V, ϕ) , and let $\text{GL}_A(V, \phi)$ be the group of units in $\text{End}_A(V, \phi)$ consisting of automorphisms of (V, ϕ) . The homomorphism $\lambda: \text{GL}_A(V, \phi) \rightarrow A^\times$ is called the *similitude character* of (V, ϕ) . We often write $\text{GL}(V, \phi)$ for $\text{GL}_{\mathbb{Q}_p}(V, \phi)$.

REMARKS.

- (a) When V is 2-dimensional, then $\text{End}_A(V, \phi) \simeq M_2(A)$ and $\lambda = \det$.
- (b) A morphism $\sigma: (V, \phi) \rightarrow (V', \phi')$ of symplectic A -modules is an isomorphism if and only if σ is bijective and $\lambda(\sigma) \in A^\times$.

DEFINITION 2.15. Let (V, ϕ) be a symplectic module over A . A representation $\rho: G \rightarrow \text{GL}_A(V)$ of a group G is *symplectic* if the image $\rho(G)$ lies in $\text{GL}_A(V, \phi)$. The associated character $\lambda \circ \rho: G \rightarrow A^\times$ is called the *similitude character* of ρ .

LEMMA 2.16. Let (V, ϕ) be a symplectic A -module. Let $\rho: G \rightarrow \text{GL}_A(V)$ be a representation, and let λ be a character of G . Then ρ is symplectic with similitude character λ if and only if given a basis of V with respect to which ϕ is represented by a matrix Q , we have

$$\rho(g)^t Q \rho(g) = \lambda(g) Q \quad \text{for all } g \in G.$$

Moreover, if ρ is symplectic with similitude character λ and $\rho^\perp := \lambda \rho^{-t}$, then $\rho^\perp \simeq \rho$.

PROOF. The proof follows by definitions. $\square$

An A -linear group representation induces an A -algebra representation of the group ring by linearity. There is a similar correspondence for symplectic group representations. We make the following definitions to make this correspondence precise.

DEFINITIONS 2.17.

- (i) Let R be an A -algebra. An *involution* on R is a bijective A -linear map $\tau: R \rightarrow R$ such that $\tau(r_1 r_2) = \tau(r_2) \tau(r_1)$ for all $r_1, r_2 \in R$. Such a tuple (R, τ) is called an *involutive algebra* over A .
- (ii) A *morphism* between involutive A -algebras (R, τ) and (R', τ') is an A -algebra map $\psi: (R, \tau) \rightarrow (R', \tau')$ such that

$$\psi(\tau(r)) = \tau'(\psi(r)) \quad \text{for all } r \in R.$$

Given a skew symmetric matrix $Q \in \text{GL}_{2n}(A)$, we define an involution τ_Q on $M_{2n}(A)$ by $\tau_Q(M) = Q^{-1} M^t Q$ for all $M \in M_{2n}(A)$.

DEFINITION 2.18. [MQ26, Definition 2.1] A *symplectic* representation of an involutive A -algebra (R, τ) is a morphism of involutive algebras $(R, \tau) \rightarrow (M_{2n}(A), \tau_Q)$, for a skew symmetric matrix Q . By a change of basis, Q can be taken to be J .

Let (V, ϕ) be a symplectic module of rank $2n$ over A , $\lambda: G \rightarrow A^\times$ be a character of G , and let $\tau = \tau_\lambda$ be the A -linear involution τ on $A[G]$ given by $\tau(g) = \lambda(g)g^{-1}$ for $g \in G$. The following result is an easy consequence of the definitions.

LEMMA 2.19. Let $\rho: G \rightarrow \text{GL}_A(V)$ be a representation and let $\tilde{\rho}: A[G] \rightarrow M_{2n}(A)$ be the induced A -algebra representation after fixing a basis of V . Let $\lambda: G \rightarrow A^\times$ be a character. Then $\tilde{\rho}$ is a symplectic representation of $(A[G], \tau_\lambda)$ if and only if ρ is a symplectic representation with similitude character λ .

2.2.1. *The general symplectic group.* Let (V, ϕ) be a symplectic module of rank $2n$ over A with a symplectic basis B (such a basis always exists by Proposition 2.5). For $\sigma \in \text{GL}_A(V, \phi)$, let M_σ be the corresponding matrix in $M_{2n}(A)$ with respect to B . The map taking $\sigma \in \text{GL}_A(V, \phi)$ to $M_\sigma \in \text{GL}_{2n}(A)$ gives an isomorphism between the group $\text{GL}_A(V, \phi)$ and the $2n$ -dimensional *general symplectic group*, denoted by $\text{GSp}_{2n}(A)$, consisting of matrices M in $\text{GL}_{2n}(A)$ satisfying

$$M^t J M = \lambda(M) J \quad \text{for some } \lambda(M) \in A^\times.$$

After fixing a symplectic basis, a symplectic representation $G \rightarrow \text{GL}_A(V, \phi)$ induces a representation $G \rightarrow \text{GSp}_{2n}(A)$.

The following result gives a precise picture of how the $(n \times n)$ blocks of a matrix in the symplectic group relate to each other. The proof of the following Lemma 2.20 follows by the definition of $\text{GSp}_{2n}(A)$ and is used implicitly in our proofs.

LEMMA 2.20. Let A be a ring with $2 \in A^\times$. A matrix $M = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix} \in \text{GL}_{2n}(A)$, where $A_i \in M_n(A)$ for $1 \leq i \leq 4$, belongs to $\text{GSp}_{2n}(A)$ if and only if

$$A_1^t A_3 \text{ and } A_2^t A_4 \text{ are symmetric; and } A_1^t A_4 - A_3^t A_2 = \lambda(M).$$

Moreover, if $M \in \text{GSp}_{2n}(A)$ and $A_3^t A_2 = 0$, then A_1 is upper (or lower) triangular if and only if A_4 is lower (or upper) triangular.

3. Symplectic types and constructing idempotents

The first part of this section is about symplectic representations and their types [MQ26, Definition 5.3]. This sets up notation to prove Proposition 1.6. In the second part, given a residually multiplicity free symplectic representation $\rho: G \rightarrow \mathrm{GSp}_{2n}(\mathbb{Z}_p)$, we construct orthogonal idempotents in $\mathbb{Z}_p[G]/\ker \rho$ corresponding to a symplectic basis (see Corollary 3.17 and Theorem 3.14). We use these idempotents in Section 4 to prove an analogue of [BG06, Proposition 3.1.1] (see Proposition 4.26).

3.1. Symmetry of symplectic representations. When working with a symplectic representation, there is a symmetry on the set of its pairwise distinct irreducible factors [BC09, §1.8]. The first part of this subsection is to explain this symmetry precisely. Later, we define symplectic types, as in [MQ26], and show that symplectic representations give rise to symplectic types.

DEFINITION 3.1. Let K be a field, and let (R, τ) be an involutive K -algebra. Let $\psi: R \rightarrow \mathrm{End}_K(V)$ be a representation. Let $V^* := \mathrm{Hom}_K(V, K)$. We define the representation $\psi^\perp: R \rightarrow \mathrm{End}_K(V^*)$ as follows: for all $r \in R$, $f \in V^*$, $v \in V$, we define

$$(1) \quad (rf)(v) = f(\tau(r)v).$$

For a representation $\psi: R \rightarrow M_n(K)$, the representation $\psi^\perp = (\psi \circ \tau)^t$

LEMMA 3.2. Let $\psi: R \rightarrow \mathrm{End}_K(V)$ be a representation and let $\psi^\perp: R \rightarrow \mathrm{End}_K(V^*)$ be the representation on V^* as defined in Definition 3.1. Then

- (a) a subspace W of V is R -stable if and only if $(V/W)^*$ is R -stable in V^* ,
- (b) a subspace of V^* is of the form $(V/W)^*$ for a subspace W of V .

In particular, $\psi^\perp$ is irreducible if and only if ψ is irreducible.

PROOF. We first prove (a). If W is an R -stable subspace of V , then $(V/W)^*$ is R -stable by Eq. (1). For the converse, if we let W' be the span of $\{\psi(r)w \mid r \in R, w \in W\}$, then for all $f \in (V/W)^*$, f vanishes on W' . Since $W \subseteq W'$ and $(V/W)^* \subseteq (V/W')^*$, using dimensions, we see that $W = W'$. Hence W is R -stable.

We now prove (b). Given a subspace $\tilde{W}$ of V^* , we let $W = \{v \in V \mid f(v) = 0 \text{ for all } f \in \tilde{W}\}$. By definition, we have $\tilde{W} \subseteq (V/W)^*$. By another dimension count argument, one can easily show that $\tilde{W} = (V/W)^*$. $\square$

The following lemma relates the Jordan–Hölder factors of $\psi^\perp$ to those of ψ .

LEMMA 3.3. Let V be a vector space over a field K , (R, τ) be an involutive K -algebra, and let $\psi: R \rightarrow \mathrm{End}_K(V)$ be a representation with $\psi^{\mathrm{ss}} \simeq \psi_1 \oplus \cdots \oplus \psi_d$, where ψ_i is irreducible for all $1 \leq i \leq d$. Then

$$(\psi^\perp)^{\mathrm{ss}} \simeq \psi_1^\perp \oplus \cdots \oplus \psi_d^\perp.$$

PROOF. When $d = 1$, the result is [Lemma 3.2](#). Assume $d \geq 2$. There is a basis of V with respect to which ψ is block upper triangular. That is, there exists $Q \in \mathrm{GL}_n(K)$ such that

$$\psi = Q \begin{pmatrix} \psi_{i_1} & * & \cdots & * \\ & \psi_{i_2} & \cdots & * \\ & & \ddots & \vdots \\ & & & \psi_{i_d} \end{pmatrix} Q^{-1}.$$

For any $r \in R$, we see that

$$\psi^\perp(r) = \psi(\tau(r))^t = Q^{-t} \begin{pmatrix} \psi_{i_1}^\perp(r) & & & \\ * & \psi_{i_2}^\perp(r) & & \\ \vdots & \vdots & \ddots & \\ * & * & \cdots & \psi_{i_d}^\perp(r) \end{pmatrix} Q^t.$$

Therefore, we get $(\psi^\perp)^{\mathrm{ss}} = \psi_1^\perp \oplus \cdots \oplus \psi_d^\perp$. $\square$

Following [\[BC09\]](#), we define the *sign* of an irreducible representation ψ such that $\psi \simeq \psi^\perp$.

LEMMA 3.4. [\[BC09, Lemma 1.8.7\]](#) *Let $\psi: R \rightarrow M_n(K)$ be an irreducible representation such that $\psi^\perp \simeq \psi$. For any $Q \in \mathrm{GL}_n(K)$ such that $\psi^\perp = Q\psi Q^{-1}$, we have $Q^t Q^{-1} = \varepsilon = \pm 1$. The sign ε does not depend on the choice of Q and is called the sign of ψ .*

PROOF. Using $\psi \simeq (\psi^\perp)^\perp$, the proof follows by Schur's lemma. $\square$

REMARK 3.5. Let (R, τ) be an involutive K -algebra, and let $\psi: R \rightarrow M_{2n}(K)$ be an irreducible representation. It is an easy calculation to see that ψ is symplectic if and only if $\psi \simeq \psi^\perp$ over K with sign -1 .

PROPOSITION 3.6. *Let $\psi: (R, \tau) \rightarrow (M_{2n}(K), \tau_Q)$ be a symplectic representation with absolutely irreducible Jordan–Hölder factors, and let $\psi_1, \dots, \psi_d$ be the isomorphism classes of the Jordan–Hölder factors of ψ . There exists a transposition σ on $I := \{1, 2, \dots, d\}$ such that $\psi_{\sigma(i)} \simeq \psi_i^\perp$ for all $i \in I$.*

PROOF. Let $Q \in \mathrm{GL}_{2n}(K)$ be a skew-symmetric matrix such that $\psi^\perp = Q\psi Q^{-1}$. Since ψ and $\psi^\perp$ are isomorphic, their semisimplifications are isomorphic. The result follows by [Lemma 3.3](#). $\square$

3.2. The symplectic extension graph is skew-symmetric. We are ready to prove [Proposition 1.6](#): the symplectic extension graph associated with an irreducible p -adic representation is a skew symmetric graph.

Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space, $\rho: G \rightarrow \mathrm{GL}(V, \phi)$ be a p -adic representation admitting a G -stable primitive lattice (that is, a symplectic $\mathbb{Z}_p$ -module of rank $\dim V$), and let $\rho_1, \dots, \rho_d$ be the distinct isomorphism classes of absolutely irreducible residual factors of ρ . By [Proposition 3.6](#), there exists a transposition σ on $\{1, \dots, d\}$ such that $\rho_{\sigma(i)} \simeq \rho_i^\perp$.

PROOF OF [PROPOSITION 1.6](#). For $1 \leq i \neq j \leq d$, assume that there is an edge (ρ_j, ρ_i) in $\Gamma_\rho^{\mathrm{GSp}}$. We need to show that $(\rho_{\sigma(i)}, \rho_{\sigma(j)})$ is an edge in $\Gamma_\rho^{\mathrm{GSp}}$. Let Λ be a G -stable symplectic vertex realizing the edge (ρ_j, ρ_i) , and let Λ_1, Λ_2 , and Λ_3 be G -stable lattices such that

$p\Lambda \subset \Lambda_1 \subset \Lambda_2 \subset \Lambda_3 \subset \Lambda$ and the subquotient $\Lambda_3/\Lambda_1 (= (\Lambda_3/p\Lambda)/(\Lambda_1/p\Lambda))$ is a nonsplit extension ρ_j by ρ_i . We have the induced G -stable flag on the dual of $\bar{\Lambda}$,

$$0 \subset \text{Hom}_{\mathbb{F}_p}(\Lambda/\Lambda_3, \mathbb{F}_p) \subset \text{Hom}_{\mathbb{F}_p}(\Lambda/\Lambda_2, \mathbb{F}_p) \subset \text{Hom}_{\mathbb{F}_p}(\Lambda/\Lambda_1, \mathbb{F}_p) \subset \text{Hom}_{\mathbb{F}_p}(\bar{\Lambda}, \mathbb{F}_p),$$

where the G -action is given by $\bar{\rho}_\Lambda^\perp$. If $\Lambda'_i := \text{Hom}_{\mathbb{F}_p}(\Lambda/\Lambda_i, \mathbb{F}_p)$ for $1 \leq i \leq 3$, then the subquotient Λ'_1/Λ'_3 of $\bar{\Lambda}^*$ is a nonsplit extension of $\rho_{\sigma(i)}$ by $\rho_{\sigma(j)}$ as Λ_3/Λ_1 is nonsplit. Since $\bar{\rho}_\Lambda \simeq \bar{\rho}_\Lambda^\perp$ by [Lemma 2.16](#), we see that Λ also realizes the edge $(\rho_{\sigma(i)}, \rho_{\sigma(j)})$ in Γ_ρ^{GSp} . $\square$

3.3. Symplectic group representations. We define symplectic (GMA) types as in [\[MQ26\]](#). Let (V, ϕ) be a symplectic vector space over a field K with $\text{Char}(K) \neq 2$. Given a symplectic multiplicity free representation $\psi: G \rightarrow \text{GL}_K(V, \phi)$, we show that it gives rise to a symplectic type.

DEFINITION 3.7. [\[MQ26, Definition 5.3\]](#) Let $I = \{1, \dots, d\}$. A *symplectic type* of dimension $2n$ is a data $\delta = ((I_0, I_1, I_2), \sigma, (n_i)_{i \in I})$, where

- (i) $\sigma: I \rightarrow I$ is a permutation such that $\sigma^2 = \text{Id}$,
- (ii) $I = I_0 \sqcup I_1 \sqcup I_2$, where $I_0 \subset I$ is the subset fixed by σ , and $\sigma(I_1) = I_2$ such that $\sigma(i) > i$ for all $i \in I_1$,
- (iii) n_i is even for all $i \in I_0$,
- (iv) $n_{\sigma(i)} = n_i$ for all $i \in I_1$,
- (v) $\sum_{i \in I} n_i = 2n$.

EXAMPLE 3.8. When $d = 2$, we have two symplectic types. Let $\delta = ((I_0, I_1, I_2), \sigma, (n_1, n_2))$ be a symplectic type of dimension $2n$. If σ is trivial, then $I_0 = \{1, 2\}$ and n_1, n_2 are even by [Definition 3.7\(iii\)](#). If σ is nontrivial, then $I_1 = \{1\}$ and $n_1 = n_2 = n$.

Let (V, ϕ) be a symplectic vector space over a field K with $\text{Char}(K) \neq 2$, and let $\psi: G \rightarrow \text{GL}_K(V, \phi)$ be a multiplicity free symplectic representation of a group G with similitude character λ . Let $\psi^{\text{ss}} \simeq \psi_1 \oplus \dots \oplus \psi_d$ and let ψ_i be absolutely irreducible for $1 \leq i \leq d$. Let $R = K[G]$ and $\tau: R \rightarrow R$ be the K -linear involution such that $\tau(g) = \lambda(g)g^{-1}$ for all $g \in G$. By [Proposition 3.6](#), there exists a transposition σ on $I = \{1, \dots, d\}$ such that $\psi_{\sigma(i)} \simeq \psi_i^\perp$. Let $I_0 = \{i \in I \mid \sigma(i) = i\}$, $I_1 = \{i \in I \mid i < \sigma(i)\}$, and let $d_1 = \#I_1$. Let n_i be the dimension of ψ_i for all $i \in I$.

PROPOSITION 3.9. *The data $\delta = ((I_0, I_1, \sigma(I_1)), \sigma, (n_i)_{i \in I})$ defines a symplectic type of dimension $2n$.*

When ψ is the residual representation associated with an irreducible symplectic representation over a local domain with residue field K , [Proposition 3.9](#) follows from [\[BC09, Lemma 1.8.8\]](#). Moreover, [Proposition 3.9](#) is a special case of [\[MQ26, Theorem 5.12\]](#). Since we are working over $\mathbb{Z}_p$, we give a simpler proof of [Proposition 3.9](#). Our proof also works for complete discrete valuation rings. The only thing left to prove is that n_i is even for all $i \in I_0$, which we prove by showing that ρ_i is symplectic for all $i \in I_0$.

PROPOSITION 3.10. *Let W be an irreducible G -stable subspace of (V, ϕ) on which the G -action is given by ψ_i for some $i \in I$. Then W is either isotropic or symplectic. Moreover, W is isotropic if and only if $i \notin I_0$.*

PROOF. Let $W' = \{v \in W \mid \phi(v, W) = 0\} = W \cap W^0$. Since W is G -stable, W' is G -stable. Moreover, W is irreducible as a G -module implies that $W' \in \{0, W\}$. This shows that W is either an isotropic or a symplectic subspace of (V, ϕ) .

Suppose that W is isotropic. By [Lemma 2.7](#), there exists a basis of V extending a basis of W such that

$$\psi = \begin{pmatrix} \psi_i & * & * \\ P_1 & P_2 \\ P_3 & P_4 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} & -I_{n_i} \\ J_{2(n-n_i)} & \\ I_{n_i} & \end{pmatrix} \psi \begin{pmatrix} & -I_{n_i} \\ J_{2(n-n_i)} & \\ I_{n_i} & \end{pmatrix}^{-1} = \psi^\perp,$$

where P_4 is a square matrix of dimension n_i , P_1 is a square matrix of dimension $2(n - n_i)$, and other block matrices have suitable dimensions. From the latter equation, we see that $P_3 \equiv 0$ on G , P_1 and P_4 are representations of G , and $P_4 = \psi_i^\perp$. Since ψ is multiplicity free, we see that $i \notin I_0$. If W is symplectic, then the result follows by [Proposition 3.11](#) below. $\square$

PROPOSITION 3.11. *Let $0 = V_0 \subset V_1 \subset \cdots \subset V_j \subset V$ be a G -stable flag such that V_{j-1} is isotropic, and V_i/V_{i-1} is irreducible with the G -action given by ψ_i for $i \in \{1, \dots, j\}$. Let $1 \leq k \leq j-1$ be the least integer such that $\phi(V_k \times V_j) \neq 0$. Then $\sigma(k) = j$.*

PROOF. Let $V'_k = \{w \in V_k \mid \phi(w, V_j) = 0\} = V_k \cap V_j^0$, which is G -stable. We see that V'_k contains V_{k-1} and $V_k \neq V'_k$ by hypothesis. Since V_k/V_{k-1} is irreducible as a G -module, we get $V'_k = V_{k-1}$. Similarly, $V'_j = V_j \cap V_k^0$ is G -stable and $V'_j = V_{j-1}$. Thus, we have shown that the induced pairing

$$\phi: (V_k/V_{k-1}) \times (V_j/V_{j-1}) \rightarrow K$$

is perfect. Therefore, ϕ induces an isomorphism between V_k/V_{k-1} and $(V_j/V_{j-1})^*$ as G -modules, where the G -action on $(V_j/V_{j-1})^*$ is given by $\psi_j^\perp$. This shows that $\psi_k \simeq \psi_j^\perp$. $\square$

LEMMA 3.12. *Assume $I_0 = I$. Then ψ is semisimple and n_i is even for all $i \in I_0$. Moreover, there is a basis of V with respect to which $\psi = \varphi_0$.*

PROOF. We induct on d . The result holds for $d = 1$ since ψ is irreducible and symplectic. Assume $d \geq 2$. Let ψ_k be a subrepresentation of ψ . Since $k \in I_0$, there is a G -stable symplectic subspace of V on which the G -action is given by ψ_k by [Proposition 3.10](#). This implies n_i is even. By [Corollary 2.10](#), there is a basis of V such that

$$\psi = \begin{pmatrix} \psi_i & * \\ & \varphi \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} J_{n_i} & \\ & J_{2n-n_i} \end{pmatrix} \psi \begin{pmatrix} J_{n_i} & \\ & J_{2n-n_i} \end{pmatrix}^{-1} = \psi^\perp.$$

The latter equation implies that $* = 0$ and $J\varphi J^{-1} = \varphi^\perp$. By induction, φ is semisimple, and n_i is even for $i \in I_0 - \{k\}$. This proves the result. $\square$

The following result shows that a maximal G -stable isotropic flag has length d_1 in V . This is the dimension of the simplicial complex of G -stable lattices in the symplectic building (see [Proposition 4.25](#)).

PROPOSITION 3.13. *Assume $I_0 \neq I$. There exists a maximal G -stable isotropic flag*

$$0 = V_0 \subset V_1 \subset \cdots \subset V_{d_1} \subset V,$$

where the G -action on V_k/V_{k-1} is given by ψ_{i_k} for all $1 \leq k \leq d_1$ and $\psi_{i_{k_1}} \not\cong \psi_{i_{k_2}}^\perp$ for all $1 \leq k_1, k_2 \leq d_1$.

PROOF. First we claim that there is an isotropic G -stable subspace of V . Let W be a G -stable subspace of V . If W is isotropic, then let $V_1 = W$. If not, then W is symplectic by Proposition 3.10. By the proof of Lemma 3.12, if the G -action on W is given by ψ_{i_0} for $i_0 \in I_0$, then there exists a subrepresentation φ of ψ such that

$$\psi^{\text{ss}} = \varphi^{\text{ss}} + \psi_{i_0}.$$

The claim now follows by inducting on d by the assumption $I_0 \neq I$.

Let

$$0 = V_0 \subset V_1 \subset \cdots \subset V_j \subset V,$$

be a maximal G -stable isotropic flag where the G -action on V_k/V_{k-1} is given by ψ_{i_k} , with $\psi_{i_{k_1}} \not\cong \psi_{i_{k_2}}^\perp$, for all $1 \leq k_1, k_2 \leq j$. Let $m = \sum_{k=1}^j n_{i_k}$. By Lemma 2.7 for the above isotropic flag, there exists a basis of V such that

$$\psi = \begin{pmatrix} \varphi & * & * \\ & P_1 & P_2 \\ & P_3 & P_4 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} & & -I_m \\ & J & \\ I_m & & \end{pmatrix} \psi \begin{pmatrix} & & -I_m \\ & J & \\ I_m & & \end{pmatrix}^{-1} = \psi^\perp.$$

Since φ is invertible, the latter equation implies that $P_3 \equiv 0$. Moreover, we see that $JP_1J^{-1} = P_1^\perp$ and $P_4 = \varphi^\perp$. If $j < d_1$, then there is a subrepresentation $\psi_{i_{j+1}}$ of P_1 such that $i_{j+1} \notin I_0$ and $\psi_{i_{j+1}} \not\cong \psi_k^\perp$ for any $1 \leq k \leq j$. That is, there exists a G -stable subspace V_{j+1} of V , containing V_j , such that G -acts on V_{j+1}/V_j by $\psi_{i_{j+1}}$. Since $i_{j+1} \notin I_0$, the subspace V_{j+1} is isotropic in V by Proposition 3.11. This contradicts the maximality of j . Thus, $j = d_1$. $\square$

PROOF OF PROPOSITION 3.9. Proposition 3.13 guarantees the existence of a maximal G -stable isotropic flag in V . Let $m = \sum_{i \in I_1} n_i$, and let $\{w_1, \dots, w_n, v_{m+1}, \dots, v_n, v_1, \dots, v_m\}$ be a basis of V as found in Lemma 2.7. With respect to such a basis,

$$\psi = \begin{pmatrix} P & * & * \\ & \varphi & * \\ & & P^\perp \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} & & -I_m \\ & J & \\ I_m & & \end{pmatrix} \psi \begin{pmatrix} & & -I_m \\ & J & \\ I_m & & \end{pmatrix}^{-1} = \psi^\perp.$$

The latter equation implies $J\varphi J^{-1} = \varphi^\perp$. Since φ is symplectic with Jordan–Hölder factors of $\{\psi_i\}_{i \in I_0}$, φ is semisimple and n_i is even for all $i \in I_0$ by Lemma 3.12. $\square$

3.4. Constructing idempotents. Let (Λ_0, ϕ) be a symplectic $\mathbb{Z}_p$ -module of rank $2n$, and let $\rho: G \rightarrow \text{GL}_{\mathbb{Z}_p}(\Lambda_0, \phi)$ be a symplectic representation that is residually multiplicity free with absolutely irreducible residual factors $\rho_1, \dots, \rho_d$. We make the following assumption in this section.

(dist) There exists $g_0 \in G$ such that $\bar{\rho}(g_0)$ has pairwise distinct eigenvalues.

In this section, we construct certain idempotents in $\mathbb{Z}_p[G]/\ker \rho$ corresponding to a symplectic basis of Λ_0 . This is used later to show that every G -stable vertex in the symplectic

building lies in an apartment (see [Proposition 4.26](#)). In the spirit of Urban [[Urb01](#)], we prove the following result to construct our idempotents.

We introduce some notation to state the result below. Let $m_0 = \sum_{i \in I_0} n_i$. Fix a diagonal embedding $\iota: \bigoplus_{i \in I_0} M_{n_i}(\mathbb{F}_p) \hookrightarrow M_{m_0}(\mathbb{F}_p)$. Let φ_0 be the composition of $\bigoplus_{i \in I_0} \rho_i$ with ι , and let $J_0 = \iota((J_{n_i})_{i \in I_0})$.

THEOREM 3.14. *Let (Λ_0, ϕ) be a symplectic $\mathbb{Z}_p$ -module of rank $2n$. Let $\rho: G \rightarrow \mathrm{GL}_{\mathbb{Z}_p}(\Lambda_0, \phi)$ be a symplectic residually multiplicity-free representation with absolutely irreducible residual factors $\rho_1, \dots, \rho_d$. Assume that ρ satisfies [\(dist\)](#). Then there exists a basis of Λ_0 with respect to which,*

- (a) $\rho(g_0)$ is diagonal,
- (b) for $m_1 = \sum_{i \in I_1} n_i$,

$$\bar{\rho} = \begin{pmatrix} P & * & * \\ & \varphi_0 & * \\ & & P^\perp \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 0 & -I_{m_1} \\ 0 & J_0 & 0 \\ I_{m_1} & 0 & 0 \end{pmatrix} \bar{\rho} \begin{pmatrix} 0 & 0 & -I_{m_1} \\ 0 & J_0 & 0 \\ I_{m_1} & 0 & 0 \end{pmatrix}^{-1} = \bar{\rho}^\perp,$$

where

$$P = \begin{pmatrix} \rho_{i_1} & * & * \\ & \ddots & * \\ & & \rho_{i_{d_1}} \end{pmatrix} \quad \text{and} \quad i_k \notin I_0.$$

REMARK 3.15. The assumption [\(dist\)](#) should be removable and should be equivalent to ρ being multiplicity free. For example, this is true when $n = 1$ using trace-determinant relations [[Bel19](#), 2.1.1]. One could ask if such a result generalizes.

We start with the following general lemma.

LEMMA 3.16. *Let (Λ_0, ϕ) be a symplectic $\mathbb{Z}_p$ -module of rank $2n$, $M \in \mathrm{GL}_{\mathbb{Z}_p}(\Lambda_0, \phi)$ with eigenvalues $\{\alpha_1, \dots, \alpha_{2n}\}$ that are distinct modulo p , and let $p\Lambda_0 \subset \Lambda_1 \subset \dots \subset \Lambda_n \subset \Lambda_0$ be a flag such that $\Lambda_i/p\Lambda_0$ is isotropic and stable under $\bar{M}$ for all $1 \leq i \leq n$. Then there exists a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ_0 such that*

- (a) $\Lambda_i/p\Lambda_0 = \langle \bar{w}_1, \dots, \bar{w}_i \rangle$ for $1 \leq i \leq n$,
- (b) $M = \mathrm{diag}(\alpha_{i_1}, \dots, \alpha_{i_{2n}})$.

PROOF. Let us denote by $\bar{V}_i$ the subspace $\Lambda_i/p\Lambda_0$ of $\bar{\Lambda}_0$ for all $1 \leq i \leq n$. We induct on $n \geq 1$. Let $n = 1$. If $\bar{V}_1 = \langle \bar{w} \rangle$, then $\bar{M}\bar{w} = \bar{\alpha}_i \bar{w}$ for some $1 \leq i \leq 2$. Let $w_1 \in \Lambda_0$ such that $Mw_1 = \alpha_i w_1$. Since $\bar{\alpha}_1 \neq \bar{\alpha}_2$, $\bar{w}_1 = a\bar{w}$ for some $a \in \mathbb{F}_p^\times$. Dividing by a , we can assume that $\bar{w}_1 = \bar{w}$. For $v_1 \in V$ such that $Mv_1 = \alpha_j v_1$ and $j \neq i$, since ϕ is nondegenerate on Λ_0 , we necessarily have $\phi(v_1, w_1) \neq 0 \pmod{p}$. By replacing v_1 by $\phi(v_1, w_1)^{-1} v_1$, we get a symplectic basis $\{w_1, v_1\}$ with respect to which the result holds.

Assume the result holds for $m < n$. Let $0 \subset \bar{V}_1 \subset \dots \subset \bar{V}_n \subset \bar{\Lambda}_0$ be an $\bar{M}$ -stable isotropic flag. Let $w_1 \in V$ such that $\bar{V}_1 = \langle \bar{w}_1 \rangle$ and $Mw_1 = \alpha_{i_1} w_1$ (the argument is same as in the previous paragraph). Let $\{w_1, w'_2, \dots, w'_n, v'_1, \dots, v'_n\}$ be a symplectic basis of $\bar{\Lambda}_0$, extending a symplectic basis of Λ_0 such that $\bar{V}_i = \langle \bar{w}_1, \bar{w}'_2, \dots, \bar{w}'_i \rangle$ for $1 < i \leq n$. With respect to the

basis $\{w_1, \dots, w'_n, v'_2, \dots, v'_n, v'_1\}$,

$$M = \begin{pmatrix} \alpha_{i_1} & *_1 & *_2 \\ & M' & *_3 \\ & *_4 & \alpha_{i_k} \end{pmatrix} \quad \text{and} \quad M^t \begin{pmatrix} & & -1 \\ & J & \\ 1 & & \end{pmatrix} M = \lambda(M) \begin{pmatrix} & & -1 \\ & J & \\ 1 & & \end{pmatrix}.$$

From the latter equation, $*_4 = 0$ and $\alpha_{i_1}\alpha_{i_k} = \lambda(M)$. The aim is to choose an eigenvector candidate for v'_1 to ensure that, for all $i \in \{1, 2, 3\}$, $*_i$ is trivial. Let $v_1 \in \Lambda_0$ be such that $Mv_1 = \alpha_{i_k}v_1$. Let $v_1 = a_1w_1 + \sum_{i=2}^n a_iw'_i + \sum_{i=1}^n b_iv'_i$. We claim that $b_1 = \phi(v_1, w_1)$ and that it is a unit in $\mathbb{Z}_p$. By our choice of v_1 , $\bar{v}_1$ has eigenvalue $\bar{\alpha}_{i_k}$. If $b_1 \equiv 0 \pmod p$, then $\bar{v}_1$ lies in a stable subspace of $\bar{\Lambda}_0$ on which the action is given by $\begin{pmatrix} \alpha_{i_1} & *_1 \\ & M' \end{pmatrix}$. This implies $\bar{\alpha}_{i_k} = \bar{\alpha}_j$ for some $j \neq i_k$, which is a contradiction. Hence, $b_1 \in \mathbb{Z}_p^\times$. Dividing by b_1 , we can assume $b_1 = \phi(v_1, w_1) = 1$. Let $W_1 = \langle w_1, v_1 \rangle$, and let $W_2 = \Lambda_0 \cap W_1^0$. Since W_1 is M -stable, we see that W_2 is M -stable. Indeed, since $MW_1 = W_1$, we have

$$\phi(MW_2, W_1) = \phi(MW_2, MM^{-1}W_1) = \lambda(M)\phi(W_2, W_1) = 0.$$

Moreover, W_2 is seen to be a symplectic $\mathbb{Z}_p$ -module of rank $2(n-1)$. Let $\tilde{\Lambda}_i = \{w - \phi(v_1, w)w_1 \mid w \in \Lambda_i\}$. Using the induction hypothesis on $M|_{W_2}$ with respect to the flag $pW_2 \subset \tilde{\Lambda}_2 \subset \dots \subset \tilde{\Lambda}_n \subset W_2$, we get a symplectic basis $\{w_2, \dots, w_n, v_2, \dots, v_n\}$ of W_2 with respect to which $\tilde{\Lambda}_i/pW_0 = \{\bar{w}_2, \dots, \bar{w}_i\}$ and $M|_{W_2}$ is diagonal. Therefore, the required basis is $\{w_1, \dots, w_n, v_1, \dots, v_n\}$. $\square$

PROOF OF THEOREM 3.14. By Hensel's lemma, $\rho(g_0)$ has eigenvalues in $\mathbb{Z}_p$, that are distinct modulo p . Let $\Lambda_1, \dots, \Lambda_{d_1}$ be $\mathbb{Z}_p$ -submodules of Λ_0 such that $p\Lambda_0 \subset \Lambda_1 \subset \dots \subset \Lambda_{d_1} \subset \Lambda_0$, and $\Lambda_k/p\Lambda_0$ is G -stable and isotropic in $\bar{\Lambda}_0$ for all $1 \leq k \leq d_1$. Let $W_0 = 0$, $W_k = \Lambda_k/p\Lambda_0$, and let the G -action on W_k/W_{k-1} be given by ρ_{i_k} for all $1 \leq k \leq d_1$. Moreover, we can extend the given G -stable isotropic flag to a G -stable flag

$$0 = W_0 \subset W_1 \subset \dots \subset W_{d_1} \subset W_{d_1+1} \subset \dots \subset W_{d_1+d_0} \subset \bar{\Lambda}_0,$$

where the G -action on W_{d_1+k}/W_{d_1+k-1} is given by ρ_{j_k} for every $k \in \{1, \dots, d_0\}$ and $j_k \in I_0$.

Since $0 \subset W_1 \subset \dots \subset W_{d_1+d_0} \subset \bar{\Lambda}_0$ is also $\bar{\rho}(g_0)$ -stable, we can consider a refined isotropic flag $0 = V_0 \subset V_1 \subset \dots \subset V_n \subset \bar{\Lambda}_0$ such that $W_k = V_{\dim W_k}$ for all $1 \leq k \leq d_1$. There is a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ_0 with respect to which $\rho(g_0)$ is diagonal with eigenvalues distinct modulo p and $V_i = \langle \bar{w}_1, \dots, \bar{w}_i \rangle$ for all $1 \leq i \leq n$ by Lemma 3.16. If $m = \dim W_k$, then the required basis is $\{w_1, \dots, w_n, v_{m+1}, \dots, v_n, v_1, \dots, v_m\}$. $\square$

Using the idempotents we construct in Corollary 3.17, we remark that Theorem 3.14 is a version of [Bel19, Proposition 2.4.2 (iii)].

COROLLARY 3.17. *Assume there exists a basis of Λ_0 satisfying the conclusions of Theorem 3.14. There exist orthogonal idempotents $\{e_i\}_{i=1}^{2n}$ in $\mathbb{Z}_p[G]/\ker \rho$, and a fixed-point-free transposition η on $\{1, \dots, 2n\}$ such that*

- (a) $e_1 + \dots + e_{2n} = 1$,
- (b) For $i \neq \eta(j)$, $(e_i + e_j)\Lambda_0$ is isotropic in Λ_0 ,
- (c) For all $i \in \{1, \dots, 2n\}$, $e_i\Lambda_0 + e_{\eta(i)}\Lambda_0$ is a symplectic submodule of Λ_0 .

Similar to [Urb01, Proof of Theorem 1.1], we need the following classical result to prove Corollary 3.17.

LEMMA 3.18. *Let $M = \text{diag}(\alpha_1, \dots, \alpha_{2n}) \in \text{GL}_{2n}(\mathbb{Z}_p)$ such that $\alpha_i \not\equiv \alpha_j \pmod{p}$ for all $1 \leq i \neq j \leq 2n$. The subalgebra of $M_{2n}(\mathbb{Z}_p)$ generated by M is the diagonal matrices.*

PROOF OF COROLLARY 3.17. Fix a basis of Λ_0 satisfying the conclusions of Theorem 3.14. By renaming the ρ_i 's, we can assume that Theorem 3.14 holds with $i_j = j$ for $1 \leq j \leq d_1$. For $1 \leq i \leq 2n$, there exists $e_i \in \mathbb{Z}_p[G]/\ker \rho$ such that $\rho(e_i) = \text{diag}(0, \dots, 0, 1, 0, \dots, 0)$, where 1 is in the i -th position, by Lemma 3.18. The transposition η on $\{1, \dots, 2n\}$ is defined such that

$$\rho(e_{\eta(i)}) = \rho^\perp(e_i).$$

By our choice of idempotents and transposition η , the result follows. $\square$

4. Buildings

A *Bruhat–Tits building* is a simplicial complex that is a union of subcomplexes called *apartments*, which are Coxeter complexes, and satisfies certain axioms [AB08, Definition 4.1]. For connected semisimple linear algebraic groups over $\mathbb{Q}_p$, the buildings constructed correspond to infinite *Weyl groups*, giving us Euclidean buildings that are CAT(0) metric spaces [AB08, Chapter 11]. In this section, we define the (Euclidean) Bruhat–Tits building associated with $\text{GL}_n(\mathbb{Q}_p)$ and $\text{GSp}_{2n}(\mathbb{Q}_p)$ whose vertices correspond to homothety class of certain $\mathbb{Z}_p$ -lattices.

Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space, and let $\rho: G \rightarrow \text{GL}(V, \phi)$ be a symplectic representation that stabilizes a primitive lattice with residual factors $\rho_1, \dots, \rho_d$ that are pairwise nonisomorphic and absolutely irreducible. The simplicial complex $\mathcal{S}_\rho^{\text{GL}}$ formed by G -stable vertices in the building associated with $\text{GL}(V)$ is compact and bounded if and only if ρ is irreducible, and has dimension $d - 1$ [Bel03, Proposition 3.2.1 and Proposition 3.3.2].

We study the subcomplex $\mathcal{S}_\rho^{\text{GSp}}$ of $\mathcal{S}_\rho^{\text{GL}}$ formed by vertices in the building associated with $\text{GL}(V, \phi)$, find its dimension (see Proposition 4.25), and, assuming ρ satisfies (dist), we show that $\mathcal{S}_\rho^{\text{GSp}}$ lies in a sub-apartment of the building associated with $\text{GL}(V, \phi)$ (see Proposition 4.26).

Suppose for all $1 \leq i \leq d$, we have $\sigma(i) \neq i$, then we can move between symplectic vertices in $\mathcal{S}_\rho^{\text{GSp}}$. Moreover, assuming ρ is irreducible, we can find symplectic vertices on the boundary (see Theorem 4.34). This is equivalent to the existence of certain edges in Γ_ρ^{GSp} (see Proposition 4.33). We use this in the proof of the main theorem Theorem 1.7.

4.1. Building associated with general linear groups. Let V be a $\mathbb{Q}_p$ -vector space and Λ be a lattice in V . The homothety class of Λ is $\{\alpha\Lambda \mid \alpha \in \mathbb{Q}_p^\times\}$ and is denoted by $[\Lambda]$. We define Bruhat–Tits buildings as a simplicial complex with homothety classes of lattices as its vertices. This is not the most natural definition, but a useful one to visualize and work with. For more equivalent definitions, see Abramenko–Brown [AB08, Chapters 4, 5, 6].

DEFINITION 4.1. Let V be an n -dimensional $\mathbb{Q}_p$ -vector space. The *Bruhat–Tits building* associated with $\text{GL}(V)$ is the following simplicial complex denoted by $\mathcal{X}_n$.

- (1) A vertex (0-simplex) is a homothety class of a lattice in V .
- (2) An edge (1-simplex) between vertices x_1, x_2 of $\mathcal{X}_n$ exists if there are representatives Λ_1 and Λ_2 for x_1 and x_2 respectively, such that

$$p\Lambda_1 \subset \Lambda_2 \subset \Lambda_1.$$

Such an x_2 called a *neighbour* of x_1 in $\mathcal{X}_n$.

- (3) An m -simplex in $\mathcal{X}_n$ is given by $(m+1)$ -vertices $x_i = [\Lambda_i]$ for all $0 \leq i \leq m$, satisfying

$$p\Lambda_0 \subset \Lambda_m \subset \Lambda_{m-1} \subset \cdots \subset \Lambda_1 \subset \Lambda_0.$$

REMARK 4.2.

- (a) The neighbour relation is symmetric: If x' is a neighbour of x , then x is also a neighbour of x' . To see this, choose representative lattices Λ and Λ' of x and x' respectively, such that

$$p\Lambda \subset \Lambda' \subset \Lambda.$$

This implies that $p(p^{-1}\Lambda') \subset \Lambda \subset p^{-1}\Lambda'$. Thus choosing the new representative $p^{-1}\Lambda'$ for x' , we see that x is a neighbour of x' .

- (b) The neighbours of a vertex $[\Lambda]$ in $\mathcal{X}_n$ are in one-to-one correspondence with the proper $\mathbb{F}_p$ -subspaces of $\Lambda/p\Lambda$. For a neighbour $[\Lambda']$ of $[\Lambda]$ such that $p\Lambda \subset \Lambda' \subset \Lambda$ the associated proper $\mathbb{F}_p$ -subspace is $\Lambda'/p\Lambda$.

4.2. Building associated with general symplectic groups. We define the Bruhat–Tits building associated with symplectic groups analogous to Definition 4.1 [Gar97, Chapter 20]. These can be seen as a subcomplex of the building $\mathcal{X}_{2n}$. We start by defining symplectic and primitive lattices in symplectic vector spaces. The vertices in the symplectic building will then be either a homothety class of a primitive vertex, or a neighbour in $\mathcal{X}_{2n}$ that corresponds under Remark 4.2(b) to an isotropic subspace.

Let (V, ϕ) be a symplectic vector space over $\mathbb{Q}_p$ of dimension $2n$. Given a basis $\{e_1, \dots, e_{2n}\}$ of V , we write $\langle e_1, \dots, e_{2n} \rangle$ for its $\mathbb{Z}_p$ -span. Let Λ be a lattice of V . There exists an integer $k \in \mathbb{Z}$ such that $\phi(\Lambda \times \Lambda) = p^k \mathbb{Z}_p$. Indeed, if $\Lambda = \langle e_1, \dots, e_{2n} \rangle$, then k is the minimum of the p -adic valuations of $\phi(e_i, e_j)$ for $1 \leq i, j \leq 2n$. Let $\bar{\Lambda} = \Lambda/p\Lambda$, and for $v \in \Lambda$, let $\bar{v}$ denote the element $v + p\Lambda$ in $\bar{\Lambda}$. Define the bilinear form $\bar{\phi}_\Lambda: \bar{\Lambda} \times \bar{\Lambda} \rightarrow \mathbb{F}_p$ by $\bar{\phi}_\Lambda(\bar{v}, \bar{w}) = \phi(v, w)p^{-k} \bmod p$, for all $v, w \in \Lambda$. Since $\phi(p\Lambda \times \Lambda)p^{-k} = \phi(\Lambda \times p\Lambda)p^{-k} = p^{k+1}p^{-k}\mathbb{Z}_p = p\mathbb{Z}_p$, the map $\bar{\phi}_\Lambda$ is well defined. Moreover, $\bar{\phi}_\Lambda$ is alternating as ϕ is alternating.

DEFINITION 4.3. A lattice Λ of V is *symplectic* if $\bar{\phi}_\Lambda$ is a symplectic pairing on $\bar{\Lambda}$. A symplectic lattice Λ is *primitive* if $\phi(\Lambda \times \Lambda) = \mathbb{Z}_p$.

As remarked in Section 2, a $\mathbb{Z}_p$ -lattice Λ in a symplectic $\mathbb{Q}_p$ -vector space (V, ϕ) is primitive if and only if $(\Lambda, \phi|_{\Lambda \times \Lambda})$ is a symplectic $\mathbb{Z}_p$ -module of rank $\dim V$. If Λ is a symplectic lattice, then the symplectic $\mathbb{F}_p$ -vector space $(\bar{\Lambda}, \bar{\phi}_\Lambda)$ only depends on the homothety class $[\Lambda]$.

PROPOSITION 4.4. Let $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ be a symplectic basis of V . The lattice

$$\Lambda = \langle p^{a_1}w_1, \dots, p^{a_n}w_n, p^{b_1}v_1, \dots, p^{b_n}v_n \rangle$$

is symplectic if and only if $a_i + b_i = a_j + b_j$ for all $1 \leq i, j \leq n$. Moreover, Λ is primitive if and only if $a_i + b_i = 0$ for all $1 \leq i \leq n$.

PROOF. Let $k = \min\{a_i + b_i \mid 1 \leq i \leq n\}$. Since $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ is a symplectic basis, we observe that $\phi(\Lambda \times \Lambda) = p^k \mathbb{Z}_p$. Moreover, for any $1 \leq i \leq n$, the element w_i pairs trivially with all the elements of the basis except v_i . If $a_i + b_i > k$ for some i , then $p^{a_i} w_i$ is such that

$$\phi(p^{a_i} w_i, p^{b_i} v_i) \equiv 0 \pmod{p^{k+1}}.$$

This implies that the alternating form $\bar{\phi}_\Lambda$ on $\bar{\Lambda}$ is degenerate. Hence, if Λ is symplectic, then $a_i + b_i = k$ for all $1 \leq i \leq n$.

For the converse, let $a_i + b_i = k$ for all $1 \leq i \leq n$. Then

$$\phi(p^{a_i} w_i, p^{b_i} v_j) = p^k \delta_{ij}, \quad \text{for all } 1 \leq i, j \leq n.$$

Since $\phi(w_i, w_j) = \phi(v_i, v_j) = 0$ and $\phi(w_i, v_j) = \delta_{ij}$ for all $1 \leq i, j \leq n$, it follows that $\bar{\phi}_\Lambda$ is given by J with respect to the basis $\{p^{a_1} w_1, \dots, p^{a_n} w_n, p^{b_1} v_1, \dots, p^{b_n} v_n\}$. This shows that Λ is symplectic. Since $\phi(\Lambda \times \Lambda) = p^k \mathbb{Z}_p$, we see that Λ is primitive if and only if $k = a_i + b_i = 0$ for all $1 \leq i \leq n$. $\square$

DEFINITION 4.5. Let (V, ϕ) be a $2n$ -dimensional symplectic $\mathbb{Q}_p$ -vector space. The *Bruhat-Tits building* associated with $\text{GL}_{\mathbb{Q}_p}(V, \phi)$ is the simplicial complex denoted by $\Delta_n(V, \phi)$ defined by the following properties.

- (1) A vertex (0-simplex) is either a homothety class of a primitive lattice, or a homothety class with a representative Λ such that there is a primitive lattice Λ_0 such that

$$p\Lambda_0 \subset \Lambda \subset \Lambda_0$$

and the subspace $\Lambda/p\Lambda_0$ is an isotropic subspace of $\bar{\Lambda}_0$.

- (2) An edge (1-simplex) between vertices x_1, x_2 in Δ_n exists if there are representatives Λ_1 and Λ_2 for x_1 and x_2 respectively, and a primitive lattice Λ_0 such that

$$p\Lambda_0 \subseteq \Lambda_1, \Lambda_2 \subseteq \Lambda_0,$$

$\Lambda_i/p\Lambda$ is isotropic for $i \in \{1, 2\}$, and either $\Lambda_1 \subset \Lambda_2$ or $\Lambda_2 \subset \Lambda_1$. Such a pair x_1, x_2 are said to be *neighbours* in Δ_n .

- (3) An m -simplex in Δ_n is given by $(m+1)$ -vertices $x_i = [\Lambda_i]$ for $0 \leq i \leq m$ where Λ_0 is a primitive lattice, and


$$p\Lambda_0 \subset \Lambda_1 \subset \Lambda_2 \subset \dots \subset \Lambda_m \subset \Lambda_0,$$

where $\Lambda_m/p\Lambda_0$ is isotropic. We denote such an m -simplex by $[x_0, \dots, x_m]$. A maximal simplex is called a *chamber*.

- (4) The *dimension* of an m -simplex is m . The *dimension* of Δ_n is the dimension of a chamber inside it.

We note that a vertex in $\Delta_n(V, \phi)$ is by definition a primitive lattice or a neighbour of a primitive lattice. For convenience, sometimes we write Δ_n for $\Delta_n(V, \phi)$.

LEMMA 4.6. *The dimension of Δ_n is n .*

FIGURE 1. 1-ball around $[0, 0, 0, 0]$.

PROOF. Let Λ_0 be a primitive lattice. Since a chamber containing $[\Lambda_0]$ in Δ_n corresponds to a maximal isotropic flag in $\bar{\Lambda}_0$, the dimension of a chamber in Δ_n is the dimension of a Lagrangian subspace of $\bar{\Lambda}_0$. Such a Lagrangian subspace has dimension n by [Lemma 2.11](#). $\square$

We note that not all vertices in Δ_n correspond to symplectic lattices (see [Example 4.8](#)). This motivates the following definition.

DEFINITION 4.7. A *symplectic (primitive) vertex* in Δ_n is the homothety class of a symplectic (primitive) lattice.

It is also common to call a primitive vertex a *hyperspecial vertex*, and a nonprimitive symplectic vertex a *special vertex*.

EXAMPLE 4.8. Let $\{w_1, w_2, v_1, v_2\}$ be a symplectic basis of $\mathbb{Q}_p^4$. For integers a_1, a_2, b_1, b_2 , let $[a_1, a_2, b_1, b_2]$ denote the homothety class $[\langle p^{a_1}w_1, p^{a_2}w_2, p^{b_1}v_1, p^{b_2}v_2 \rangle]$. The shape of the 1-ball around $[0, 0, 0, 0]$ in Δ_2 is seen in [Fig. 1](#). Each dashed line represents an edge between symplectic vertices.

4.2.1. *Symplectic vertices and their neighbours.* Let (V, ϕ) be a symplectic vector space over $\mathbb{Q}_p$ of dimension $2n$. Let x be a symplectic vertex in Δ_n and let y be its neighbour. By definition, there exists representative lattices Λ and Λ' of x and y respectively, and a primitive lattice Λ_0 such that $p\Lambda_0 \subset \Lambda', \Lambda \subset \Lambda_0$ with $\Lambda' \subset \Lambda$ or $\Lambda \subset \Lambda'$. We will see in this section that $\Lambda/p\Lambda_0$ Lagrangian subspace of $\Lambda_0/p\Lambda_0$ (see [Corollary 4.13](#)). Therefore, we must have $p\Lambda \subset \Lambda' \subset \Lambda$. Moreover, one can show that $\Lambda'/p\Lambda$ is isotropic in $\Lambda/p\Lambda$ (for example, using [Proposition 4.12](#)).

We start with some basic properties of primitive lattices in a symplectic vector space, and show that primitive lattices are symplectic $\mathbb{Z}_p$ -modules (see [Corollary 4.11](#)).

PROPOSITION 4.9. *Let Λ be a primitive lattice of V . Let $w_1, v_1 \in \Lambda$ such that $\phi(v_1, w_1) = 1$, $\Lambda_1 = \langle w_1, v_1 \rangle$ be the $\mathbb{Z}_p$ -span of $\{w_1, v_1\}$, and let $\Lambda_2 = \Lambda_1^0 \cap \Lambda$. Then $\Lambda = \Lambda_1 \oplus \Lambda_2$. Moreover, Λ_2 is a primitive lattice of $\Lambda_2 \otimes \mathbb{Q}_p$ with respect to the symplectic form induced by ϕ .*

PROOF. Since $\phi(\Lambda \times \Lambda) = \mathbb{Z}_p$, for any $w \in \Lambda$, the element $\tilde{w} = w + \phi(w, v_1)w_1 - \phi(w, w_1)v_1$ lies in Λ_2 . Hence, $\Lambda = \Lambda_1 + \Lambda_2$. Moreover, $\Lambda_2 \cap \Lambda_1 = 0$ since Λ_1 is a symplectic submodule of Λ . This shows that $\Lambda = \Lambda_1 \oplus \Lambda_2$.

We claim that the symplectic vector space $\bar{\Lambda}$ decomposes as

$$\bar{\Lambda} = (\Lambda_1 + p\Lambda)/p\Lambda \oplus (\Lambda_2 + p\Lambda)/p\Lambda.$$

Indeed, if $v = v_1 + v_2$, for $v_1 \in \Lambda_1$ and $v_2 \in \Lambda_2$, then $\bar{v} = \bar{v}_1 + \bar{v}_2 = (\bar{v}_1 - p\bar{w}) + (\bar{v}_2 + p\bar{w})$. This proves that the two spaces span $\bar{\Lambda}$. To prove the above direct sum decomposition, we show that $(\Lambda_1 + p\Lambda) \cap (\Lambda_2 + p\Lambda) = p\Lambda$. Let $v_1 \in \Lambda_1$ and $v_2 \in \Lambda_2$ such that $v_1 + pv = v_2 + pw$ lies in the intersection $(\Lambda_1 + p\Lambda) \cap (\Lambda_2 + p\Lambda)$. Since $\Lambda = \Lambda_1 \oplus \Lambda_2$, there exists $w_1 \in \Lambda_1$ and $w_2 \in \Lambda_2$ such that $v_1 - v_2 = p(w - v) = pw_1 - pw_2$. Since $\Lambda_1 \cap \Lambda_2 = 0$, we see that $v_i = pw_i$ for $i \in \{1, 2\}$. This proves the claim.

We show that Λ_2 is a primitive lattice. Since $\phi(v_1, w_1) = 1$, we see that $(\Lambda_1 + p\Lambda)/p\Lambda$ is a symplectic subspace of $\bar{\Lambda}$. We see that $(\Lambda_2 + p\Lambda)/p\Lambda$ is also symplectic by [Corollary 2.9](#). If $\phi(\Lambda_2 \times \Lambda_2) \subseteq p\mathbb{Z}_p$, then $\bar{\phi}_\Lambda \equiv 0$ on $(\Lambda_2 + p\Lambda)/p\Lambda$. This proves that $\phi(\Lambda_2 \times \Lambda_2) = \mathbb{Z}_p$. We only need to show $\bar{\Lambda}_2 = \Lambda_2/p\Lambda_2$ is a symplectic vector space. Let $w \in \Lambda_2$ such that $\phi(w, \Lambda_2) \subseteq p\mathbb{Z}_p$. Since $\Lambda_2 + p\Lambda = \Lambda_2 + p\Lambda_1$ and $\phi(\Lambda_2 \times \Lambda_1) = 0$, we have $\phi(w, \Lambda_2 + p\Lambda) \subseteq p\mathbb{Z}_p$. Thus, the element $w \in p\Lambda$ as $\bar{\Lambda}$ is symplectic. Let $w_2 \in \Lambda_2$ such that $w - pw_2 \in p\Lambda_1$. Since $w, w_2 \in \Lambda_2$ and $\Lambda_2 \cap \Lambda_1 = 0$, we have $w = pw_2 \in p\Lambda_2$. This proves the result. $\square$

LEMMA 4.10. *Let (V, ϕ) be a $2n$ -dimensional symplectic $\mathbb{Q}_p$ -vector space and let Λ_0 be a primitive lattice in V . A symplectic basis of $\bar{\Lambda}_0$ can be lifted to a symplectic basis of Λ_0 .*

PROOF. We induct on n . By [Proposition 2.5](#), the residual symplectic vector space $\bar{\Lambda}_0$ has a symplectic basis; say $\{\bar{w}_1, \dots, \bar{w}_n, \bar{v}_1, \dots, \bar{v}_n\}$. Let w_1 and v_1 be a lift of $\bar{w}_1$ and $\bar{v}_1$, respectively, in Λ_0 . Since $\phi(v_1, w_1) \equiv 1 \pmod{p}$, by replacing v_1 by $\phi(v_1, w_1)^{-1}v_1$, we can assume that $\phi(v_1, w_1) = 1$. Let $\Lambda_1 = \langle w_1, v_1 \rangle$, and let $\Lambda_2 = \Lambda_0 \cap \Lambda_1^0$. The lattice Λ_2 is primitive in $\Lambda_2 \otimes \mathbb{Q}_p$ by [Proposition 4.9](#). Moreover, for a lift $w \in \Lambda_0$ of $\bar{w} \in \{\bar{w}_i, \bar{v}_i\}_{i=2}^n$, we see that $\tilde{w}$ lies in Λ_2 and lifts $\bar{w}$, as $\phi(w, v_1) \equiv \phi(w, w_1) \equiv 0 \pmod{p}$. Therefore, there exist lifts of elements of $\{\bar{w}_i, \bar{v}_i\}_{i=2}^n$ in Λ_2 . Since the dimension of $\Lambda_2 \otimes \mathbb{Q}_p$ is $2(n-1)$, by the induction hypothesis, there is a symplectic basis $\{w_2, \dots, w_n, v_2, \dots, v_n\}$ of Λ_2 lifting the symplectic basis $\{\bar{w}_2, \dots, \bar{w}_n, \bar{v}_2, \dots, \bar{v}_n\}$. Therefore, the symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ for Λ_0 lifts our given symplectic basis of $\bar{\Lambda}_0$. $\square$

COROLLARY 4.11. *A $\mathbb{Z}_p$ -lattice in (V, ϕ) is primitive if and only if it is spanned by a symplectic basis. In particular, a primitive $\mathbb{Z}_p$ -lattice Λ is a symplectic $\mathbb{Z}_p$ -module with respect to $\phi|_{\Lambda \times \Lambda}$.*

The following result ensures that every m -simplex in Δ_n is contained in an apartment (see [Definition 4.18](#)) of Δ_n .

PROPOSITION 4.12. *Let Λ_0 be a primitive lattice in V and let $\Lambda_1, \dots, \Lambda_m$ be lattices such that $p\Lambda_0 \subset \Lambda_1 \subset \dots \subset \Lambda_m \subset \Lambda_0$ and the subspace $\Lambda_m/p\Lambda_0$ is isotropic in $\bar{\Lambda}_0$. For all $1 \leq i \leq m$, let $k_i = \dim_{\mathbb{F}_p}(\Lambda_i/p\Lambda_0)$. There exists a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ_0 such that*

$$\Lambda_i = \langle w_1, \dots, w_{k_i}, pw_{k_i+1}, \dots, pw_n, pv_1, \dots, pv_n \rangle.$$

PROOF. Let $V_i = \Lambda_i/p\Lambda_0$ for $1 \leq i \leq m$. By Proposition 2.12, there is a symplectic basis $\{\bar{w}_1, \dots, \bar{w}_n, \bar{v}_1, \dots, \bar{v}_n\}$ of $\bar{\Lambda}_0$ such that $\{\bar{w}_1, \dots, \bar{w}_{k_i}\}$ spans V_i for all $1 \leq i \leq m$. There is a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ_0 such that w_i lifts $\bar{w}_i$ for all $1 \leq i \leq n$ by Lemma 4.10. For all $1 \leq i \leq m$, let

$$V'_i = \langle w_1, \dots, w_{k_i}, pw_{k_i+1}, \dots, pw_n, pv_1, \dots, pv_n \rangle/p\Lambda_0.$$

Since $\dim_{\mathbb{F}_p} V'_i = k_i$ and $V'_i \subseteq V_i$, the result follows. $\square$

The following result characterizes symplectic neighbours of a primitive vertex in Δ_n .

COROLLARY 4.13. *Let Λ_0 be a primitive lattice in V and let Λ be a lattice such that $p\Lambda_0 \subset \Lambda \subset \Lambda_0$ and $\Lambda/p\Lambda_0$ is an isotropic subspace of $\bar{\Lambda}_0$. The lattice Λ is symplectic if and only if $\Lambda/p\Lambda_0$ is Lagrangian.*

PROOF. By Proposition 4.12, there exists a symplectic basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ_0 such that

$$\Lambda = \langle w_1, \dots, w_s, pw_{s+1}, \dots, pw_n, pv_1, \dots, pv_n \rangle.$$

It follows that Λ is symplectic if and only if $s = n$ by Proposition 4.4. $\square$

COROLLARY 4.14. *A chamber in Δ_n contains exactly one primitive vertex and exactly one special vertex.*

PROOF. Let $[x_0, \dots, x_n]$ be a chamber. Choose representative lattices Λ_i for x_i for all $i \in \{0, \dots, n\}$ such that Λ_0 is primitive, and $p\Lambda_0 \subset \Lambda_1 \subset \dots \subset \Lambda_n \subset \Lambda_0$. This implies that $\Lambda_n/p\Lambda_0$ is Lagrangian in $\bar{\Lambda}_0$. The lattice Λ_n must be symplectic with $\phi(\Lambda_n \times \Lambda_n) = p\mathbb{Z}_p$ by Corollary 4.13. $\square$

COROLLARY 4.15. *Let x, x' be neighbouring symplectic vertices in Δ_n . Then one of x or x' is represented by a primitive lattice.*

PROOF. Let Λ and Λ' be representatives of x and x' respectively, and Λ_0 be a primitive lattice such that $p\Lambda_0 \subseteq \Lambda' \subset \Lambda \subseteq \Lambda_0$ and $\Lambda/p\Lambda_0$ is isotropic. Then by Corollary 4.13, either $\Lambda' = p\Lambda_0$ or $\Lambda = \Lambda_0$. $\square$

PROPOSITION 4.16. *Let x be a special vertex in Δ_n . There exists a primitive vertex $x_0 \in \Delta_n$ such that x and x_0 are neighbours in Δ_n .*

PROOF. Since x is a special vertex, we can find a representative lattice Λ for x such that $\phi(\Lambda \times \Lambda) = p\mathbb{Z}_p$. Since x is symplectic, there is a $\mathbb{Z}_p$ -basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ that lifts a symplectic basis of $\bar{\Lambda}$; that is, we have $\phi(v_i, w_i) = p$ for all $1 \leq i \leq n$, $\phi(v_i, w_j) \equiv 0 \pmod{p^2}$ for all $1 \leq i \neq j \leq n$, and $\phi(v_i, v_j) \equiv \phi(w_i, w_j) \equiv 0 \pmod{p^2}$ for all $1 \leq i, j \leq n$. Let Λ_0 be the

$\mathbb{Z}_p$ -span of $\{w_1, \dots, w_n, p^{-1}v_1, \dots, p^{-1}v_n\}$. We see that $p\Lambda_0 \subset \Lambda \subset \Lambda_0$ and $\phi(\Lambda_0 \times \Lambda_0) = \mathbb{Z}_p$ since $\phi(w_1, p^{-1}v_1) = 1$. We only need to show $\bar{\Lambda}_0$ is symplectic. Let $w_0 \in \Lambda_0$ such that $\phi(w_0, w) \equiv 0 \pmod p$ for all $w \in \Lambda_0$. If $w_0 = \sum_{i=1}^n a_i w_i + \sum_{i=1}^n b_i p^{-1}v_i$, then

$$a_i \equiv \phi(w_0, p^{-1}v_i) \equiv 0 \pmod p \quad \text{and} \quad b_i \equiv \phi(w_0, w_i) = 0 \pmod p,$$

for all $1 \leq i \leq n$. This shows that $w_0 \in p\Lambda_0$. $\square$

We can characterize symplectic neighbours of symplectic vertices in Δ_n by [Proposition 4.16](#) and [Corollary 4.13](#).

COROLLARY 4.17. *Let $x = [\Lambda]$ be a symplectic vertex such that $\phi(\Lambda \times \Lambda) \in \{\mathbb{Z}_p, p\mathbb{Z}_p\}$. Let $y = [\Lambda']$ be a neighbour of x in Δ_n such that $p\Lambda \subset \Lambda' \subset \Lambda$. Then y is symplectic if and only if $\Lambda'/p\Lambda$ is Lagrangian in $\bar{\Lambda}$.*

PROOF. If Λ is primitive, then the result is the same as [Corollary 4.13](#). Let Λ be symplectic with $\phi(\Lambda \times \Lambda) = p\mathbb{Z}_p$. Let B be a symplectic basis of $\bar{\Lambda}$ extending a basis of $\Lambda'/p\Lambda$. We can lift B to a basis $\{w_1, \dots, w_n, v_1, \dots, v_n\}$ of Λ . By the same argument as in the proof of [Proposition 4.16](#), the lattice $\Lambda_0 = \langle p^{-1}w_1, \dots, p^{-1}w_n, v_1, \dots, v_n \rangle$ is primitive. Moreover, we see that $p\Lambda \subset \Lambda' \subseteq p\Lambda_0 \subset \Lambda \subset \Lambda_0$. Since $p\Lambda_0/p\Lambda$ is spanned by the images of $w_1, \dots, w_n$ in $\bar{\Lambda}$, it is Lagrangian in $\bar{\Lambda}$. Since $p\Lambda_0 \subset p^{-1}\Lambda' \subseteq \Lambda_0$, the vertex y is symplectic if and only if $\Lambda' = p\Lambda_0$ by [Corollary 4.13](#). Thus, y is symplectic if and only if $\Lambda'/p\Lambda$ is Lagrangian in $\bar{\Lambda}$. $\square$

4.2.2. Apartments and sub-apartments in the symplectic building. A building is a union of apartments, which are much simpler to work with. Crucial to our proof of the main theorem is showing that every G -stable vertex in Δ_n lies in a single apartment (see [Proposition 4.26](#)).

Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space over of dimension $2n$. Using [[Set08](#), Lemma 1.3.5], we define apartments in the building Δ_n .

DEFINITION 4.18. An *apartment* in $\Delta_n = \Delta_n(V, \phi)$ corresponding to a symplectic basis $B = \{w_1, \dots, w_n, v_1, \dots, v_n\}$ of (V, ϕ) is the simplicial complex $\Sigma = \Sigma(B)$ formed by vertices in Δ_n of the form

$$[\langle p^{a_1}w_1, \dots, p^{a_n}w_n, p^{b_1}v_1, \dots, p^{b_n}v_n \rangle],$$

where $a_i, b_i \in \mathbb{Z}$ for all $1 \leq i \leq n$. We denote such a vertex in Σ by $[a_1, \dots, a_n, b_1, \dots, b_n]$.

REMARK 4.19. Let Σ be an apartment in Δ_n corresponding to a symplectic basis of V . Let $x_0 = [0, \dots, 0]$ be the origin of Σ . For all $0 \leq i \leq n$, we let $x_i = [0, \dots, 0, 1, \dots, 1]$ where the number of zeros is i . We call the chamber $[x_0, x_1, \dots, x_n]$ around x_0 as the *fundamental chamber* in Σ . Translating the fundamental chamber by the action of the affine Weyl group of $\mathrm{GSp}_{2n}(\mathbb{Q}_p)$, we obtain Σ . For an example, see [[She07](#), Example 2.3].

The following result characterizes vertices in an apartment of Δ_n .

LEMMA 4.20. [[Set08](#), Proposition 1.3.4] *Let Σ be an apartment in Δ_n corresponding to a symplectic basis of V . A vertex $y = [a_1, \dots, a_n, b_1, \dots, b_n] \in \Sigma$ if and only if there exists $m \in \mathbb{Z}$ such that*

$$a_i + b_i + 2m \in \{1, 2\}, \quad \text{for all } 1 \leq i \leq n.$$

We define sub-apartments of Δ_n analogous to [Bel03, §2.2]. Let $\mathcal{P} = \{B_j\}_{j=1}^d$ be a partition of $\{1, \dots, 2n\}$, and let

$$\mathbb{Z}_{\mathcal{P}}^{2n} = \{(a_1, \dots, a_{2n}) \in \mathbb{Z}^{2n} \mid a_i = a_{i'} \text{ for any } i, i' \in B_j\}.$$

We define a bijective mapping

$$(2) \quad \mathbb{Z}_{\mathcal{P}}^{2n} \rightarrow \mathbb{Z}^d, \quad (a_1, \dots, a_{2n}) \mapsto (b_1, \dots, b_d),$$

where $a_i = b_j$ for all $i \in B_j$.

DEFINITION 4.21. A *sub-apartment* of an apartment Σ in Δ_n is a subcomplex that under Eq. (2) corresponds to an apartment in Δ_m for some $m \leq n$. The dimension of the sub-apartment in this case is m .

We say a subcomplex Σ' of Σ is *contained in an apartment* of Δ_m if Σ' is contained in a sub-apartment of dimension m .

Let B be a fixed symplectic basis of V , and let Σ be the corresponding apartment in Δ_n . Let $\Sigma_{\mathcal{P}}$ be the subcomplex of Σ whose vertices have representative in $\mathbb{Z}_{\mathcal{P}}^{2n}$. Under further assumptions on $\mathcal{P}$, we show that $\Sigma_{\mathcal{P}}$ is a sub-apartment of Σ in Δ_n .

LEMMA 4.22. Let $B = \{w_1, \dots, w_n, v_1, \dots, v_n\}$ be a symplectic basis of V , and let

$$(\epsilon_1, \dots, \epsilon_{2n}) := (w_1, \dots, w_n, v_1, \dots, v_n).$$

Let $I = \{1, \dots, d\}$, and let $\mathcal{P} = \{B_j\}_{j \in I}$ be a partition of $\{1, \dots, 2n\}$ such that

- (1) there is an order 2 permutation σ on I ;
- (2) for all $k \neq \sigma(j)$, the subspace spanned by $\{\epsilon_i\}_{i \in B_j \cup B_k}$ is isotropic;
- (3) for all $j \in I$, the subspace spanned by $\{\epsilon_i\}_{i \in B_j \cup B_{\sigma(j)}}$ is symplectic.

Let $I_0 = \{j \in I \mid j = \sigma(j)\}$, $I_1 = \{j \in I \mid j < \sigma(j)\}$, and let $d_s = \#I_s$ for all $s \in \{0, 1\}$. Then the subcomplex $\Sigma_{\mathcal{P}}$ is a sub-apartment of the Σ of dimension $d_0 + d_1$.

PROOF. Since $B_j \cup B_{\sigma(j)}$ is symplectic, $i \in B_j$ if and only if $n+i \in B_{\sigma(j)}$. For each $j \in I_0 \sqcup I_1$, there exists $1 \leq i_j \leq n$ such that $\{\epsilon_{i_j}, \epsilon_{n+i_j}\}$ spans a symplectic subspace of V . Let V' be the symplectic subspace of V generated by $B' = \{\epsilon_{i_1}, \dots, \epsilon_{i_{d_0+d_1}}, \epsilon_{n+i_1}, \dots, \epsilon_{n+i_{d_0+d_1}}\}$. By definition, $\Sigma_{\mathcal{P}}$ is the subcomplex of Σ whose vertices are of the form $[a_1, \dots, a_{2n}]$ with $a_i = b_j$ for all $i \in B_j$. Let us define the map

$$\mathbb{Z}_{\mathcal{P}}^{2n} \rightarrow \mathbb{Z}^{2(d_0+d_1)}, \quad [a_1, \dots, a_{2n}] \mapsto [b_1, \dots, b_{2(d_0+d_1)}].$$

We see that $b_j + b_{\sigma(j)} = a_i + a_{n+i}$ for any $i \in B_j$. We see that $[b_1, \dots, b_{2(d_0+d_1)}] \in \Delta_{d_0+d_1}$ if and only if $[a_1, \dots, a_{2n}] \in \Delta_n$ by Lemma 4.20. This shows that $\Sigma_{\mathcal{P}}$ is a sub-apartment of Σ of dimension $d_0 + d_1$. $\square$

REMARK 4.23. We elaborate on the hypothesis of Lemma 4.22 by providing an example where it holds. Let $\rho: G \rightarrow \mathrm{GL}(V\phi)$ be a symplectic p -adic representation that stabilizes a G -stable primitive lattice, and with residual factors $\rho_1, \dots, \rho_d$ that are absolutely irreducible and pairwise nonisomorphic. We show later that there exists a symplectic basis B of V with respect to which every G -stable lattice in Δ_n lies in the apartment $\Sigma(B)$ and is of the form $[a_1, \dots, a_{2n}]$ (see Proposition 4.26). Moreover, if $\Lambda_0 = [0, \dots, 0]$, then the residual

representation $\bar{\rho}_{\Lambda_0}$ is block-upper triangular with respect to a basis as in [Theorem 3.14](#). Since ρ_j is irreducible for all $j \in I$, there exists a partition $\mathcal{P} = \{B_j\}_{j \in I}$ of $\{1, \dots, 2n\}$, satisfying the hypothesis of [Lemma 4.22](#), such that for all $j \in I$ and for all $i, k \in B_j$, we have $a_i = a_k$.

4.3. Simplicial complex formed by stable vertices. Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space of dimension $2n$, and let Δ_n denote the Bruhat–Tits building associated with $\mathrm{GL}(V, \phi)$. Let $\rho : G \rightarrow \mathrm{GL}(V, \phi)$ be a symplectic representation such that V has a G -stable primitive lattice. Assume ρ is residually multiplicity free, and with residual factors $\rho_1, \dots, \rho_d$ that are absolutely irreducible. Let $I = \{1, \dots, d\}$, and let n_i be the dimension of ρ_i for all $i \in I$.

DEFINITION 4.24. Let S_ρ^{GSp} be the set of G -stable vertices in Δ_n . Let $\mathcal{S}_\rho^{\mathrm{GSp}}$ be the simplicial subcomplex formed by these vertices in Δ_n . The *dimension* of $\mathcal{S}_\rho^{\mathrm{GSp}}$ is defined to be the dimension of the maximal simplex contained in $\mathcal{S}_\rho^{\mathrm{GSp}}$.

Let σ be the transposition on I such that $\rho_{\sigma(i)} \simeq \rho_i^\perp$, $I_0 = \{i \in I \mid \sigma(i) = i\}$, $I_1 = \{i \in I \mid i < \sigma(i)\}$, and let $d_1 = \#I_1$.

PROPOSITION 4.25. *The dimension of $\mathcal{S}_\rho^{\mathrm{GSp}}$ is d_1 .*

PROOF. Let Λ_0 be a G -stable primitive lattice. If $d_1 = 0$, then $\bar{\rho}_{\Lambda_0}$ only has symplectic G -stable subspaces. This means that $[\Lambda_0]$ does not have any neighbours in S_ρ^{GSp} . This implies that $\dim \mathcal{S}_\rho^{\mathrm{GSp}} = 0 = d_1$.

Let $d_1 \geq 1$. There exists a maximal G -stable isotropic flag in $\bar{\Lambda}_0$ by [Proposition 3.13](#). That is, there exists lattices $\Lambda_1, \dots, \Lambda_{d_1}$ such that

$$p\Lambda_0 \subset \Lambda_1 \subset \dots \subset \Lambda_{d_1} \subset \Lambda_0,$$

such that $\Lambda_{d_1}/p\Lambda_0$ is isotropic in $\bar{\Lambda}_0$. This proves the result. $\square$

4.3.1. Contained in an apartment. The idempotents constructed in [\[BG06\]](#) shows that $\mathcal{S}_\rho^{\mathrm{GL}}$ is contained in a $(d-1)$ -dimensional apartment in $\mathcal{X}_d$. We use [Corollary 3.17](#) to prove the analogous result for $\mathcal{S}_\rho^{\mathrm{GSp}}$.

PROPOSITION 4.26. *Let $\rho : G \rightarrow \mathrm{GL}(V, \phi)$ be a residually multiplicity free symplectic representation admitting a G -stable primitive lattice. If there exists a basis of V satisfying the conclusions of [Theorem 3.14](#), then $\mathcal{S}_\rho^{\mathrm{GSp}}$ is contained in an apartment of $\Delta_{d_0+d_1}$.*

PROOF. The strategy is the same as [\[BG06, Proposition 3.1.1\]](#). Let Λ_0 be a G -stable primitive lattice. By assumption, there exists a basis B of Λ_0 that satisfies the conclusions of [Theorem 3.14](#). By [Corollary 3.17](#), there exist orthogonal idempotents $\{e_i\}_{i=1}^{2n} \subset \mathbb{Z}_p[G]/\ker \rho$, and a transposition η such that (a), (b), and (c) of [Corollary 3.17](#) hold. We see that $\Lambda_0 = \bigoplus_{i=1}^{2n} e_i \Lambda_0$ by [Corollary 3.17 \(b\)](#). By [\[BG06, Lemma 3.1.3\]](#), any G -stable lattice $\Lambda \subset \Lambda_0$ is of the form $\Lambda = \bigoplus_{i=1}^{2n} p^{a_i} (e_i \Lambda_0)$ for some integers $a_1, \dots, a_{2n}$. Let B_0 be a symplectic basis of Λ_0 formed by permuting elements of B . By (b) and (c) of [Corollary 3.17](#), there exists a symplectic basis of Λ_0 such that any G -stable lattice $\Lambda \subset \Lambda_0$ is given by $[a_1, \dots, a_{2n}]$,

for some integers $a_1, \dots, a_{2n}$. This shows that $\mathcal{S}_\rho^{\text{GSp}}$ is contained in the apartment of Δ_n corresponding to B_0 .

To prove $\mathcal{S}_\rho^{\text{GSp}}$ is contained in a sub-apartment of dimension $d_0 + d_1$, we use [Lemma 4.22](#). Let $\mathcal{P} = \{B_j\}_{j=1}^d$ be the partition of $\{1, \dots, 2n\}$ such that the idempotent $\mathbf{e}_j = \sum_{i \in B_j} e_i$ corresponds to the residual factor ρ_j in the following sense: the map $R \rightarrow \mathbf{e}_j R \mathbf{e}_j \xrightarrow{\bar{\rho}_{\Lambda_0}} M_{2n}(\mathbb{F}_p)$ is given by

$$r \mapsto \begin{pmatrix} 0 & & & \\ & \ddots & & \\ & & \rho_j(r) & \\ & & & \ddots \\ & & & & 0 \end{pmatrix}.$$

With respect to the permutation σ on I such that $\rho_{\sigma(j)} \simeq \rho_j^\perp$, the hypothesis of [Lemma 4.22](#) is satisfied, and the result follows. $\square$

4.3.2. Moving between stable symplectic lattices. In this section, we show the equivalence of the necessary assumption $I_0 = \emptyset$ (for [Theorem 1.7](#)) and the existence of G -stable symplectic neighbours. This is also crucial to our method of proof.

Let $\rho : G \rightarrow \text{GL}(V, \phi)$ be a symplectic representation such that V has a G -stable primitive lattice. Suppose that ρ is residually multiplicity free and has residual factors $\rho_1, \dots, \rho_d$ that are absolutely irreducible. Let $I = \{1, \dots, d\}$ and $I_0 = \{i \in I \mid \sigma(i) = i\}$.

PROPOSITION 4.27. *There exists G -stable symplectic neighbours in Δ_n if and only if $I_0 = \emptyset$.*

PROOF. Let $[\Lambda]$ and $[\Lambda_0]$ be G -stable symplectic neighbours in Δ_n . Without loss of generality, we can assume that Λ_0 is primitive and $p\Lambda_0 \subset \Lambda \subset \Lambda_0$ by [Corollary 4.15](#). By [Corollary 4.13](#), $\Lambda/p\Lambda_0$ is a Lagrangian subspace of $\bar{\Lambda}_0$. By [Lemma 2.11](#), there exists a basis of $\bar{\Lambda}_0$ such that

$$\bar{\rho}_{\Lambda_0} = \begin{pmatrix} \psi & * \\ & \psi' \end{pmatrix} \quad \text{and} \quad J\bar{\rho}_{\Lambda_0}J^{-1} = \bar{\rho}_\Lambda^\perp.$$

This implies that $\psi' = \psi^\perp$, and that $I_0 = \emptyset$, since $\bar{\rho}_{\Lambda_0}$ has distinct residual factors.

We prove the converse by using [Proposition 4.25](#). If $I_0 = \emptyset$, then $d = 2d_1$ is even and the dimension of $\mathcal{S}_\rho^{\text{GSp}} = d_1$. For a G -stable primitive lattice Λ_0 , there exists an isotropic G -stable flag

$$0 = V_0 \subset V_1 \subset \dots \subset V_{d_1} \subset \bar{\Lambda}_0.$$

We see that V_{d_1} is a Lagrangian subspace of $\bar{\Lambda}_0$ since

$$2 \dim V_{d_1} = 2 \sum_{i=1}^{d_1} \dim(V_i/V_{i-1}) = 2 \sum_{i=1}^{d_1} n_i = \sum_{i=1}^{d_1} n_i + \sum_{i=1}^{d_1} n_{\sigma(i)} = 2n.$$


The G -stable Lagrangian subspace V_{d_1} of $\bar{\Lambda}_0$ corresponds to a G -stable symplectic neighbour of Λ_0 by [Corollary 4.13](#). $\square$

4.3.3. Jordan–Hölder colouring. Let $\rho: G \rightarrow \mathrm{GL}(V)$ be a symplectic p -adic representation that is residually multiplicity-free with absolutely irreducible factors $\rho_1, \dots, \rho_d$. Let $\mathcal{S}_\rho^{\mathrm{GL}}$ be the subcomplex formed by G -stable vertices in the building $\mathcal{X}_{2n}$ associated with $\mathrm{GL}(V)$. We define the Jordan–Hölder colouring on the edges in $\mathcal{S}_\rho^{\mathrm{GL}}$ as in [Bel03]. For a representation $\varphi: G \rightarrow \mathrm{GL}_n(\mathbb{F}_p)$, we let $\mathrm{JH}(\varphi)$ be the sum of the irreducible factors of φ (with multiplicities) viewed as an element of the Grothendieck group of $\mathbb{F}_p[G]$.

DEFINITION 4.28. [Bel03, §3.4.2] Let $x = [\Lambda] \in S_\rho^{\mathrm{GL}}$, and let $y = [\Lambda']$ be a neighbour of x in S_ρ^{GL} such that $p\Lambda \subset \Lambda' \subset \Lambda$. We let $\mathrm{JH}(x, y) := \mathrm{JH}(\Lambda'/p\Lambda)$.

For $x_1, x_2 \in S_\rho^{\mathrm{GL}}$ that are neighbours in $\mathcal{X}_{2n}$, we have $\mathrm{JH}(x_2, x_1) + \mathrm{JH}(x_1, x_2) = \sum_{i=1}^d \rho_i$ [Bel03, Proposition 3.4.3 (c)]. Suppose now that $\rho: G \rightarrow \mathrm{GL}(V, \phi)$ is symplectic and residually multiplicity-free. The above colouring on the edges in $\mathcal{S}_\rho^{\mathrm{GL}}$ restricts to a colouring on the edges of $\mathcal{S}_\rho^{\mathrm{GSp}}$. For a symplectic vertex $x \in S_\rho^{\mathrm{GSp}}$, and $y \in S_\rho^{\mathrm{GSp}}$ neighbour of x in Δ_n , there exists $\{\varepsilon_i\}_{i=1}^d \subset \{0, 1\}$ such that $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$ for all $i \in I$, and $\mathrm{JH}(x, y) = \sum_{i=1}^d \varepsilon_i \rho_i$. The condition $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$ is equivalent to the subspace corresponding to y being isotropic.

EXAMPLE 4.29. If $I_0 = \emptyset$, then we have 2^{d_1} -colourings of edges between G -stable symplectic vertices. When $d_1 = 1$, we have the following two colourings:



We end this section with a crucial result of [Bel03] that will be used in the next section to prove the existence of symplectic vertices in $\mathcal{S}_\rho^{\mathrm{GSp}}$ with no edges of a specific colour. Before we state the result, we define when two edges are parallel in an apartment. Let Σ be an apartment with respect to a basis in $\mathcal{X}_{2n}$, and let $x_i = [a_1^{(i)}, \dots, a_{2n}^{(i)}]$ for $1 \leq i \leq 4$. The edge (x_1, x_2) is *parallel* to (x_3, x_4) if $a_j^{(2)} - a_j^{(1)} = a_j^{(4)} - a_j^{(3)}$ for all $1 \leq j \leq 2n$.

PROPOSITION 4.30. [Bel03, Propositions 3.4.3 (a) and 3.4.4]. *Let Σ be an apartment in $\mathcal{X}_{2n}$ containing $\mathcal{S}_\rho^{\mathrm{GL}}$. Let $x, y, x', y' \in S_\rho^{\mathrm{GL}}$ such that x, y and x', y' are neighbours in $\mathcal{X}_{2n}$. Then $\mathrm{JH}(x, y) = \mathrm{JH}(x', y')$ if and only if the edges (x, y) and (x', y') are parallel in Σ .*

4.3.4. Neighbours and extensions. In this section, we show the existence of certain edges in the symplectic extension graph associated with an irreducible symplectic p -adic representation. This is the main ingredient to our proof of Theorem 1.7.

PROPOSITION 4.31. *Let $\rho: G \rightarrow \mathrm{GL}(V, \phi)$ be an irreducible symplectic p -adic representation that admits a G -stable primitive lattice and with absolutely irreducible residual factors $\rho_1, \dots, \rho_d$. Let $I = \{1, \dots, d\}$ and let σ be the transposition on I defined by $\rho_{\sigma(i)} \simeq \rho_i^\perp$ for all $i \in I$. Assume ρ satisfies (rmf) and (dist), and that $\sigma(i) \neq i$ for all $i \in I$. Given positive integers ε_i for all $i \in I$ such that $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$, there exists a symplectic vertex $x = [\Lambda] \in S_\rho^{\mathrm{GSp}}$ such that $\Lambda/p\Lambda$ admits a nonsplit extension of ρ_i by ρ_j , satisfying $\varepsilon_i = 1$ and $\varepsilon_j = 0$, as a subquotient.*

We start with a result on the existence of triangles in $\mathcal{S}_\rho^{\text{GSp}}$ with two symplectic vertices.

LEMMA 4.32. *Let $\rho: G \rightarrow \text{GL}(V, \phi)$ be a p -adic symplectic representation that admits a G -stable primitive lattice, satisfies (rmf), and let $\rho_1, \dots, \rho_d$ be its absolutely irreducible residual factors. Let $I = \{1, \dots, d\}$ and assume that $\sigma(i) \neq i$ for all $i \in I$. Let $x \in S_\rho^{\text{GSp}}$ be a symplectic vertex and let $y \in S_\rho^{\text{GL}}$ be a neighbour of x such that $\text{JH}(x, y) = \sum_{i \in I} \varepsilon_i \rho_i$. Then $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$ for all $i \in I$ if and only if y is a neighbour of x in Δ_n . If y is a neighbour of x in Δ_n , then there exists a symplectic vertex $z \in S_\rho^{\text{GSp}}$ neighbouring x and y such that $\text{JH}(x, z) = \text{JH}(x, y) + \text{JH}(y, z)$.*

PROOF. Let $x = [\Lambda]$. We see that y corresponds to a G -stable isotropic subspace $\Lambda'/p\Lambda$ of $\Lambda/p\Lambda$ if and only if $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$ for all $i \in I$ by Proposition 3.11. If $\Lambda'/p\Lambda$ is isotropic, then it is contained in a G -stable Lagrangian subspace $\Lambda''/p\Lambda$ of $\Lambda/p\Lambda$ by Proposition 4.25. This Lagrangian subspace of $\Lambda/p\Lambda$ corresponds to a symplectic neighbour z of x by Corollary 4.17. Since $p\Lambda \subset \Lambda' \subset \Lambda'' \subset \Lambda$, by definition it follows that $\text{JH}(x, z) = \text{JH}(x, y) + \text{JH}(y, z)$. $\square$

The nonexistence of G -stable neighbours in $\mathcal{S}_\rho^{\text{GL}}$ with certain JH-colouring is equivalent to the existence of certain edges in Γ_ρ^{GL} by [Bel03, Proposition 3.4.5]. We have the following analogue in our setting:

PROPOSITION 4.33. *Let $x \in S_\rho^{\text{GSp}}$ and let Λ be a symplectic lattice such that $x = [\Lambda]$. For all $i \in I$, let $\varepsilon_i \geq 0$ be an integer such that $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$. The following statements are equivalent:*

- (a) *There does not exist a neighbour $y \in S_\rho^{\text{GSp}}$ of x such that $\text{JH}(x, y) = \sum_i \varepsilon_i \rho_i$.*
- (b) *$\Lambda/p\Lambda$ admits a nonsplit extension of some ρ_i by ρ_j with $\varepsilon_i = 1$ and $\varepsilon_j = 0$.*

PROOF. Suppose (a) is false. Then there exists neighbour y of x in S_ρ^{GSp} such that $\text{JH}(x, y) = \sum_{i \in I} \varepsilon_i \rho_i$. Since y is also a neighbour of x in S_ρ^{GL} , (b) is false by [Bel03, Proposition 3.4.5]. This shows (b) implies (a). Suppose (b) is false. Then every extension of ρ_i by ρ_j with $\varepsilon_i = 1$ and $\varepsilon_j = 0$ that is a subquotient of $\Lambda/p\Lambda$ is split. There exists a neighbour $y \in S_\rho^{\text{GL}}$ of x such that $\text{JH}(x, y) = \sum_{i \in I} \varepsilon_i \rho_i$ by [Bel03, Proposition 3.4.5]. Since $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$, the G -stable subspace of $\bar{\Lambda}$ corresponding to y is isotropic by Lemma 4.32. That is $y \in S_\rho^{\text{GSp}}$. This implies that (a) implies (b). $\square$

The final ingredient for proving Proposition 4.31 is the following result that guarantees the existence of symplectic vertices with no edges of a specific colour.

THEOREM 4.34. *Assume that ρ is irreducible, satisfies (rmf) and (dist), and that $I_0 = \emptyset$. For all $i \in I$, let ε_i be a positive integer such that $\varepsilon_i + \varepsilon_{\sigma(i)} \leq 1$. There exists a symplectic vertex $x \in S_\rho^{\text{GSp}}$ with no neighbours $y \in S_\rho^{\text{GSp}}$ with $\text{JH}(x, y) = \sum_{i \in I} \varepsilon_i \rho_i$.*

PROOF. Without loss of generality, we assume $\varphi = \rho_1$. By Proposition 4.26, $\mathcal{S}_\rho^{\text{GSp}}$ lies inside an apartment Σ , corresponding to a symplectic basis B , of Δ_{d_1} . That is, every vertex in $\mathcal{S}_\rho^{\text{GSp}}$ is of the form $[a_1, \dots, a_n, b_1, \dots, b_n]$. Let $x_0 = [0, \dots, 0] \in S_\rho^{\text{GSp}}$ be the origin of Σ .

We prove by contradiction. Let us assume every symplectic vertex $x \in S_\rho^{\text{GSp}}$ has a neighbour $y \in S_\rho^{\text{GSp}}$ such that $\text{JH}(x, y) = \rho_1$. Let y_0 be the neighbour of x_0 such that $\text{JH}(x_0, y_0) = \rho_1$. Since ρ_1 is irreducible, we see that $y_0 = [1, \dots, 1, 0, 1, \dots, 1]$ where all but one entry is 1. Without loss of generality, by permuting the elements of B , we assume $y_0 = [0, 1, \dots, 1]$. Since ρ is irreducible, S_ρ^{GSp} is finite by [Bel03, Proposition 3.2.1]. Thus, there exists an integer μ , maximal such that

$$x_\mu = [0, a_2, \dots, a_{d_1}, \mu, b_2, \dots, b_{d_1}] \in S_\rho^{\text{GSp}}$$

is symplectic. By our assumption, there exists a neighbour $y_\mu \in S_\rho^{\text{GSp}}$ of x_μ such that $\text{JH}(x_\mu, y_\mu) = \rho_1$. By [Bel03, Proposition 3.4.4], since $y_0 = [0, 1, \dots, 1]$, we should have

$$y_\mu = [0, a_2 + 1, \dots, a_{d_1} + 1, \mu + 1, b_2 + 1, \dots, b_{d_1} + 1].$$

By Lemma 4.32, there exists a symplectic vertex $z \in S_\rho^{\text{GSp}}$, neighbouring x_0 , such that

$$z = [0, \dots, \mu + 1, \dots].$$

This contradicts maximality of μ . □

PROOF OF PROPOSITION 4.31. Follows by Theorem 4.34 and Proposition 4.33. □

5. Symplectic extension graph

In this section, we prove our main results: Theorem 1.7 and Theorem 1.8. We end by discussing an example where the symplectic extension graph is an empty graph. Throughout, our setup is as follows. Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space, and let $\rho: G \rightarrow \text{GL}(V, \phi)$ be a symplectic representation that admits a G -stable primitive lattice.

REMARK 5.1. An important class of representations satisfying our hypothesis are continuous p -adic irreducible symplectic representations of a compact group G . Indeed, from the Bruhat–Tits fixed point theorem [AB08, Theorem 11.23], and [AB08, Theorem 11.34], the maximal compact subgroups of $\text{GSp}_{2n}(\mathbb{Q}_p)$ are stabilizers of vertices in the building and are conjugate over $\text{GL}_{2n}(\mathbb{Q}_p)$ to $\text{GSp}_{2n}(\mathbb{Z}_p)$. Therefore, by conjugating ρ if necessary, we can assume that ρ admits a G -stable primitive lattice.

5.1. Proof of Theorem 1.7. We prove our main theorem in parts. Let $\rho_1, \dots, \rho_d$ be the absolutely irreducible classes of residual factors of ρ . Let $I = \{1, \dots, d\}$, $\sigma: I \rightarrow I$ be defined such that $\rho_{\sigma(i)} \simeq \rho_i^\perp$, and let $I_0 = \{i \in I \mid \sigma(i) = i\}$.

PROPOSITION 5.2 (THEOREM 1.7(a)). *Assume ρ is irreducible, satisfies (rmf) and (dist), and $I_0 = \emptyset$. For every $i \in I$, there is a path from ρ_i to $\rho_{\sigma(i)}$ in Γ_ρ^{GSp} .*


PROOF. Let $i \in I$ and let $\Psi := \{\psi_1, \dots, \psi_r\}$ be the vertices in Γ_ρ^{GSp} to which there is a directed path from ρ_i . Let $\psi_0 = \rho_i$ and let $\varphi = \bigoplus_{i=0}^r \psi_i$. If $k = \sigma(j)$ for some $0 \leq j \neq k \leq r$, then we are done by Proposition 1.6. Indeed, if there is a path from ρ_i to ρ_j and $\rho_{\sigma(j)}$, then there is a path from ρ_i to $\rho_{\sigma(i)}$. If $\sigma(j) \neq k$ for any $0 \leq j, k \leq r$, then there exists a $0 \leq j \leq r$ and an edge from ψ_j to ψ' in Γ_ρ^{GSp} , with $\psi' \notin \Psi$ by Proposition 4.31. A path from ρ_i to ψ_j exists in Γ_ρ^{GSp} and gives a path from ρ_i to ψ' . This is a contradiction. □

THEOREM 5.3 (**THEOREM 1.7(b)**). *Assume ρ is irreducible, satisfies (rmf) and (dist), and $I_0 = \emptyset$. The directed graph Γ_ρ^{GSp} has at most d_1 strongly connected components. Moreover, when Γ_ρ^{GSp} has d_1 strongly connected components, then $\Gamma_\rho^{\text{GSp}} = \bigsqcup_{i=1}^{d_1} \Gamma_i$, where Γ_i is the connected directed graph on ρ_i and $\rho_{\sigma(i)}$.*

PROOF. Let ψ_0 be a vertex of Γ_ρ^{GSp} . Let Γ_{ψ_0} be the subgraph of Γ_ρ^{GSp} whose vertices ψ are either equal to ψ_0 or $\psi_0^\perp$, or has a path to and from ψ_0 and $\psi_0^\perp$. We claim that

- (i) $\Gamma_{\psi_0} = \Gamma_{\psi_0^\perp}$ and it is a strongly connected subgraph of Γ_ρ^{GSp} ;
- (ii) $\psi \in \Gamma_{\psi_0}$ if and only if $\psi^\perp \in \Gamma_{\psi_0}$.

We note that (ii) follows by **Proposition 1.6**. Let us prove (i). By definition, it is immediate that $\Gamma_{\psi_0} = \Gamma_{\psi_0^\perp}$. If ψ_1 and ψ_2 belong to Γ_{ψ_0} , then we see that there is a path from ψ_1 to ψ_2 :



In (3), the dashed line represents a path. This proves (i).

We claim that there is no edge from Γ_ψ to $\Gamma_{\psi'}$ for $\psi \notin \Gamma_{\psi'}$. If there is an edge from Γ_ψ to $\Gamma_{\psi'}$, then there is a path from ψ to ψ' . There is a path from $\psi'^\perp$ to $\psi^\perp$ by **Proposition 1.6**. Thus, we see that ψ has path to and from ψ' and $\psi'^\perp$ by **Proposition 5.2**. That is, $\psi \in \Gamma_{\psi'}$. This shows that Γ_ρ^{GSp} is the disjoint union of strongly connected subgraphs Γ_ψ .

Since Γ_ψ contains the two vertices $\{\psi, \psi^\perp\}$, the maximum number of strongly connected components is $d_1 = d/2$. Moreover, Γ_ρ^{GSp} has d_1 connected components if and only if Γ_ψ is the strongly connected graph on $\{\psi, \psi^\perp\}$. $\square$

5.2. Two Jordan–Hölder factors. Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space, and let $\rho : G \rightarrow \text{GL}(V, \phi)$ be a symplectic representation admitting a G -stable primitive lattice. When ρ has two residual factors, then we have a full characterization of Γ_ρ^{GSp} .

THEOREM 5.4. *Assume ρ is irreducible, and satisfies (dist) and (rmf). The symplectic extension graph Γ_ρ^{GSp} associated with ρ is strongly connected if and only if $I_0 = \emptyset$.*

The *if* part follows from **Theorem 1.7** when $d = 2$. The *only if* part follows from the result below.

PROPOSITION 5.5. *Let $I_0 = \{1, 2\}$, and let $\text{JH}(\bar{\rho}^{\text{ss}}) = \rho_1 + \rho_2$. For any symplectic G -stable lattice Λ , $\bar{\rho}_\Lambda$ is semisimple.*

PROOF. By **Proposition 1.6**, if $[\Lambda] \in S_\rho^{\text{GSp}}$ is symplectic such that $\bar{\rho}_\Lambda$ is a nonsplit extension of ρ_2 by ρ_1 , then it is a nonsplit extension of ρ_1 by ρ_2 . $\square$

EXAMPLE 5.6. We provide an example where the symplectic extension graph is disconnected and a proper subgraph of Bellaïche's connected extension graph.

Let G be the elements in $\mathrm{GSp}_4(\mathbb{Z}_p)$ whose reduction modulo p is the set of matrices $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ with A, B, C, D are diagonal modulo p ; that is,

$$G \bmod p = \mathrm{GL}_2(\mathbb{F}_p) \times_{\mathrm{GL}_1(\mathbb{F}_p)} \mathrm{GL}_2(\mathbb{F}_p) \subset \mathrm{GSp}_4(\mathbb{F}_p).$$

More precisely,

$$G = \left\{ \begin{pmatrix} a_1 & pa_2 & b_1 & pb_2 \\ pa_3 & a_4 & pb_3 & b_4 \\ c_1 & pc_2 & d_1 & pd_2 \\ pc_3 & c_4 & pd_3 & d_4 \end{pmatrix} \mid a_1d_1 - b_1c_1 \equiv a_4d_4 - b_4c_4 \bmod p \right\}.$$

Let the representation $\rho : G \hookrightarrow \mathrm{GSp}_4(\mathbb{Z}_p)$ be the inclusion map. Then ρ is irreducible, and $\bar{\rho} = \rho_1 \oplus \rho_2$, where $\rho_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \bmod p$ is absolutely irreducible, for $1 \leq i \leq 2$, and $\det \rho_1 = \det \rho_2$. To show $\rho_1 \not\cong \rho_2$, we show that $\mathrm{Tr} \rho_1 \neq \mathrm{Tr} \rho_2$. Indeed, for

$$g = \begin{pmatrix} 1 & & & \\ & & & -1 \\ & & 1 & \\ & 1 & & \end{pmatrix},$$

we see that $\mathrm{Tr} \rho_1(g) = 2$ and $\mathrm{Tr} \rho_2(g) = 0$. Hence, the irreducible representation ρ is residually reducible and multiplicity free. Moreover, $\rho_1 \rho_2^t$ is not a character as $(\rho_1 \rho_2^t)(g)$ is not a scalar. By Proposition 5.5, $\Gamma_\rho^{\mathrm{GSp}}$ is disconnected.

5.3. Four Jordan–Hölder factors. Let (V, ϕ) be a symplectic $\mathbb{Q}_p$ -vector space, and let $\rho : G \rightarrow \mathrm{GL}(V, \phi)$ be a symplectic representation admitting a G -stable symplectic lattice and has residual factors ρ_1, ρ_2, ρ_3 , and ρ_4 that are absolutely irreducible and pairwise nonisomorphic. Let $I = \{1, 2, 3, 4\}$ and let $I_0 = \{i \in I \mid \sigma(i) = i\}$.

THEOREM 5.7 (THEOREM 5.3: $d_1 = 2$). *Assume ρ is irreducible, satisfies (dist) and (rmf), and assume that $I_0 = \emptyset$. The directed graph $\Gamma_\rho^{\mathrm{GSp}}$ is either strongly connected, or $\Gamma_\rho^{\mathrm{GSp}} = \Gamma_1 \sqcup \Gamma_2$, where Γ_i is the strongly connected graph on $\{\rho_i, \rho_i^\perp\}$ for $1 \leq i \leq 2$.*

EXAMPLE 5.8. We illustrate examples for the two cases of Theorem 5.7: one where the symplectic extension graph is strongly connected and one where it is a union of two strongly connected subgraphs. We generalize this example in the next section to show that Theorem 1.7 is optimal (see Proposition 5.13).

We first provide an example where the symplectic extension graph is strongly connected. Let

$$G_1 = \left\{ \begin{pmatrix} a_{11} & pa_{12} & a_{13} & pa_{14} \\ pa_{21} & a_{22} & pa_{23} & a_{24} \\ pa_{31} & pa_{32} & a_{33} & pa_{34} \\ pa_{41} & pa_{42} & pa_{43} & a_{44} \end{pmatrix} \in \mathrm{GSp}_4(\mathbb{Z}_p) \mid a_{ij} \in \mathbb{Z}_p \text{ for all } 1 \leq i, j \leq 4 \right\}.$$

Let $\rho: G_1 \hookrightarrow \mathrm{GSp}_4(\mathbb{Z}_p)$ be the inclusion. It is not hard to see that ρ is irreducible (or equivalently, $\mathbb{Q}_p[G_1] = M_4(\mathbb{Q}_p)$; see [Lemma 5.12](#)) and residually reducible with four Jordan–Hölder factors:

$$\rho_1 := a_{11} \bmod p, \rho_2 := a_{22} \bmod p \quad \text{and} \quad \rho_1^\perp := a_{33} \bmod p, \rho_2^\perp := a_{44} \bmod p.$$

The diagonal matrices in $\mathrm{GSp}_4(\mathbb{Z}_p)$ lie in G_1 . Let B be the symplectic basis fixed. Since B satisfies the conclusions of [Theorem 3.14](#), $\mathcal{S}_\rho^{\mathrm{GSp}}$ lies in the apartment Σ associated to B of Δ_2 . Let $x_0 = [0, \dots, 0]$ be the origin of Σ . By construction, x_0 is G_1 -stable and primitive.

Any symplectic vertex x in Σ is of the form $x = [\alpha_1, \alpha_2, \beta_1, \beta_2]$ with $\alpha_1 + \beta_1 = \alpha_2 + \beta_2$ by [Proposition 4.4](#). If x is a symplectic G_1 -stable vertex, then for every $g \in G_1$, we have

$$\mathrm{diag}(p^{\alpha_1}, p^{\alpha_2}, p^{\beta_1}, p^{\beta_2})\rho(g)\mathrm{diag}(p^{-\alpha_1}, p^{-\alpha_2}, p^{-\beta_1}, p^{-\beta_2}) \in \mathrm{GSp}_4(\mathbb{Z}_p).$$

Since a_{31}, a_{42} can be chosen such that they do not lie in $p\mathbb{Z}_p$, we get

$$0 \leq \alpha_1 - \beta_1 \leq 1 \quad \text{and} \quad 0 \leq \alpha_2 - \beta_2 \leq 1.$$


We claim either $x = x_0$ or $x = x_1 := [1, 1, 0, 0]$.

We see that $\alpha_1 = \beta_1$ if and only if $\alpha_2 = \beta_2$. If $\alpha_i = \beta_i$ and $\alpha_j = \beta_j + 1$ for $j \neq i$, then


$$2\alpha_i = \alpha_i + \beta_i = \alpha_j + \beta_j = 2\alpha_j + 1,$$

a contradiction. If $\alpha_i = \beta_i$ for some $i \in \{1, 2\}$, then $\alpha_1 = \beta_1 = \alpha_2 = \beta_2$. Therefore, $x = x_0$. If $\alpha_1 = 1 + \beta_1$, then $\alpha_2 = 1 + \beta_2$. Using $\alpha_1 + \beta_1 = \alpha_2 + \beta_2$, we see that $\alpha_1 = \alpha_2 = 1$ and $\beta_1 = \beta_2 = 0$. This shows that $x = x_1$.

Therefore, x_0 has a unique G_1 -stable neighbour $x_1 = [1, 1, 0, 0]$. The vertex x_0 realizes a nonsplit extension of $\rho_1^\perp$ by ρ_1 , and a nonsplit extension of $\rho_2^\perp$ by ρ_2 . Moreover, x_1 realizes a nonsplit extension of ρ_1 by $\rho_1^\perp$ and $\rho_2^\perp$, and ρ_2 by $\rho_2^\perp$ and $\rho_1^\perp$. The following diagram depicts $\mathcal{S}_\rho^{\mathrm{GSp}}$ in Σ :



In the above diagram, $\circ$ denotes a neighbouring vertex in $\mathcal{S}_\rho^{\mathrm{GSp}}$ of the symplectic vertices x_0 and x_1 . In this case, the symplectic extension graph $\Gamma_\rho^{\mathrm{GSp}}$ is



We provide an example where $\Gamma_\rho^{\mathrm{GSp}}$ has two strongly connected components. Let

$$G_2 = \left\{ \begin{pmatrix} a_{11} & pa_{12} & a_{13} & pa_{14} \\ pa_{21} & a_{22} & pa_{23} & a_{24} \\ pa_{31} & p^2a_{32} & a_{33} & pa_{34} \\ p^2a_{41} & pa_{42} & pa_{43} & a_{44} \end{pmatrix} \in G \right\}.$$

We see that G_2 is a subgroup of G . Let $\rho' := \rho|_{G_2}$, $\rho'_i := \rho_i|_{G_2}$ and let $\rho'^{\perp}_i := \rho_i^{\perp}|_{G_2}$ for all $1 \leq i \leq 2$. Then ρ' is an irreducible representation with residual factors $\rho'_1, \rho'_2, \rho'^{\perp}_1$, and $\rho'^{\perp}_2$. By the same calculations as above, there exist two G_2 -stable symplectic vertices x_0 and x_1 , and by the shape of G_2 , the edges from ρ_2 to $\rho_1^{\perp}$, and ρ_1 to $\rho_2^{\perp}$ are not realized by $\bar{\rho}'_{x_1}$. In this case, the symplectic extension graph $\Gamma_{\rho'}^{\text{GSp}}$ is

$$\begin{array}{cc} \rho'_1 & \rho'_2 \\ \updownarrow & \updownarrow \\ \rho'^{\perp}_1 & \rho'^{\perp}_2 \end{array}$$

We note that although ρ and ρ' have the same image modulo p , their symplectic extension graphs are different.

EXAMPLE 5.9. Let $\mathbb{F}_p(r)$ be the $G_{\mathbb{Q}}$ -module $\mathbb{F}_p$ with the $G_{\mathbb{Q}}$ -action given by multiplication by the r -th power of the mod- p cyclotomic character. Let E_k denote the normalized Eisenstein series of weight $k \geq 4$ level $\text{SL}_2(\mathbb{Z})$, and let f be a cuspidal eigenform of weight k and level $\text{SL}_2(\mathbb{Z})$ such that $f \equiv E_k \pmod{p}$. If ρ is an irreducible symplectic 4-dimensional Galois representation associated to a genus 2 cuspidal Siegel eigenform that is congruent to the Klingen Eisenstein series associated to f , then one of the two following situations must occur by [Theorem 5.7](#) and [Proposition 1.6](#):

- (a) $H^1(G_{\mathbb{Q}}, \mathbb{F}_p(1-k)) \neq 0 \neq H^1(G_{\mathbb{Q}}, \mathbb{F}_p(k-1))$.
- (b) $H^1(G_{\mathbb{Q}}, \mathbb{F}_p(2-k)) \neq 0 \neq H^1(G_{\mathbb{Q}}, \mathbb{F}_p(k-2))$.

5.4. Proof of Theorem 1.8. In this section, we prove [Theorem 1.8](#) by generalizing [Example 5.8](#) (see [Proposition 5.13](#) below).

Let $n \geq 1$ be an integer, and let σ be the transposition on $I_{2n} = \{1, \dots, 2n\}$ defined by $\sigma(i) = i + n$ for $1 \leq i \leq n$. Let r be an integer fixed throughout this section such that $1 \leq r \leq n$. Let $\{\eta_{i,j}\}_{1 \leq i,j \leq n}$ satisfy the following conditions.

- (1) For all $1 \leq i \leq n$, let $\eta_{i,i} = \eta_{i,\sigma(i)} = \eta_{\sigma(i),\sigma(i)} = \eta_{\sigma(i),i} - 1 = 0$.
- (2) For all $1 \leq i \neq j \leq n$, let $\eta_{i,j} = \eta_{i,\sigma(j)} = \eta_{\sigma(i),\sigma(j)} = 1$.
- (3) For all $1 \leq i \leq r-1$, let $\eta_{\sigma(i),j} = 2 = \eta_{\sigma(j),i}$ for every $1 \leq j \leq n$ with $j \neq i$; else $\eta_{\sigma(i),j} = \eta_{\sigma(j),i} = 1$.

Let us denote the assignment $(i, j) \mapsto \eta_{i,j}$ defined above by $\eta^{(r)}$.

DEFINITIONS 5.10. [[BG06](#), 2.1.1 and 2.1.2] A *directed numbered graph* is a triple (V, E, η) , where (V, E) is a directed graph with vertex set V and edge set $E \subseteq V^2$, and $\eta : E \rightarrow \mathbb{Z} \sqcup \{\infty\}$ is a map. We say (V, E, η) is a *metric graph* if $E = V^2$, $\eta(v, v) = 0$ for all $v \in V$, and $\eta(u, v) \leq \eta(u, w) + \eta(w, v)$ for all $u, v, w \in V$.

LEMMA 5.11. *The directed numbered graph $(I_{2n}, I_{2n}^2, \eta^{(r)})$ is a metric graph.*

PROOF. We need to show for all $i, j, k \in I_{2n}$, $\eta_{i,j} \leq \eta_{i,k} + \eta_{k,j}$. The inequality is true for $i = j$ by definition. Suppose that $i \neq j$. We see that

$$\eta_{i,k} + \eta_{k,j} = \begin{cases} \eta_{i,j} & \text{if } k \in \{i, j\}, \\ \eta_{\sigma(i),j} & \text{if } k \in \{\sigma(i), \sigma(j)\}, \\ \geq 2 & \text{otherwise.} \end{cases}$$

Since $\eta_{i,j} = \eta_{\sigma(i),j} \leq 2$, the result follows. $\square$

Let G_r be the collection of elements $(a_{ij}) \in \mathrm{GSp}_{2n}(\mathbb{Z}_p)$ with $a_{ij} \in p^{n_{i,j}}\mathbb{Z}_p$. We have the inclusions

$$G_n \subset \cdots \subset G_2 \subset G_1.$$

LEMMA 5.12. *For any $1 \leq r \leq n$, we have $\mathbb{Q}_p[G_r] = M_{2n}(\mathbb{Q}_p)$.*

PROOF. Let E_{ij} be the matrix in $M_{2n}(\mathbb{Q}_p)$ with 1 at the (i, j) -th entry and 0 everywhere else. We show that $E_{ij} \in \mathbb{Q}_p[G_r]$ for all $1 \leq i, j \leq 2n$. Since there exists a diagonal element $g_0 \in G_r$ with residually distinct eigenvalues, $E_{i,i} \in \mathbb{Q}_p[G_r]$ for all $1 \leq i \leq 2n$ by Lemma 3.18. Let σ be the transposition on $\{1, \dots, 2n\}$ such that $\sigma(k) = k + n$ for all $1 \leq k \leq 2n$. Let $\mathrm{Id} \in M_{2n}(\mathbb{Z}_p)$ be the identity matrix. By a simple calculation, for all $1 \leq i \leq 2n$, we see that $p^{\eta_{i,\sigma(i)}}E_{i,\sigma(i)} + \mathrm{Id} \in G_r$. This implies that $E_{i,\sigma(i)} \in \mathbb{Q}_p[G_r]$ for all $1 \leq i \leq 2n$. Let $1 \leq i, j \leq 2n$ be such that $j \notin \{i, \sigma(i)\}$. It remains to show that $E_{i,j} \in \mathbb{Q}_p[G_r]$. Since $\eta_{i,j} = \eta_{\sigma(j),\sigma(i)}$, we see that $\mathrm{Id} + p^{\eta_{i,j}}(E_{i,j} - E_{\sigma(j),\sigma(i)}) \in G_r$. This implies that $E_{i,j} - E_{\sigma(j),\sigma(i)} \in \mathbb{Q}_p[G_r]$. Since $E_{i,i} \in \mathbb{Q}_p[G_r]$ and $E_{i,j} = E_{ii}(E_{i,j} - E_{\sigma(j),\sigma(i)})$, we see that $E_{i,j} \in \mathbb{Q}_p[G_r]$. $\square$

PROPOSITION 5.13. *The inclusion $\rho: G_r \hookrightarrow \mathrm{GSp}_{2n}(\mathbb{Z}_p)$ is an irreducible representation whose residual factors are nonisomorphic characters. The symplectic extension graph $\Gamma_\rho^{\mathrm{GSp}}$ has r connected components.*

PROOF. By Lemma 5.11 and [BG06, Lemme 3.2.1], G_r is a subgroup of $\mathrm{GSp}_{2n}(\mathbb{Z}_p)$. Moreover, the map $\rho: G_r \hookrightarrow \mathrm{GSp}_{2n}(\mathbb{Z}_p)$ defined by inclusion is irreducible by Lemma 5.12.

For all $1 \leq i \leq 2n$, the residual factors are

$$\chi_i((a_{jk})) = a_{ii} \bmod p.$$

By definition, the residual characters $\chi_1, \dots, \chi_{2n}$ are pairwise distinct. Moreover, if $\sigma(i) = i + 1$ for $1 \leq i \leq n$, then we see that $\chi_{\sigma(i)} = \chi_i^\perp$ for all $i \in I_{2n}$. That is, if $\lambda: G_r \rightarrow \mathbb{Z}_p^\times$ is the similitude character of ρ , then $\chi_{\sigma(i)}\chi_i = \bar{\lambda}$.

Let B_0 be the symplectic basis fixed of $\mathbb{Z}_p^{2n}$. Since the diagonal matrices of $\mathrm{GSp}_{2n}(\mathbb{Z}_p)$ are in G_r , $\mathcal{S}_\rho^{\mathrm{GSp}}$ lies in the apartment of Δ_n associated to B_0 by Proposition 4.26. If $x = [\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n]$ is a G -stable symplectic vertex, then we see that $0 \leq \alpha_i - \beta_i \leq 1$ for all $i \in \{1, \dots, 2n\}$. This follows by the condition $\eta_{\sigma(i),i} = 1$ for all $1 \leq i \leq n$. By a computation similar to the one done in Example 5.8, we see the only G -stable symplectic vertices are $x_0 = [\Lambda_0] = [0, \dots, 0]$, and $x_1 = [1, \dots, 1, 0, \dots, 0]$. By our definition of $\eta^{(r)}$, in particular (3), the graph $\Gamma_\rho^{\mathrm{GSp}}$ is a disjoint union of r strongly connected subgraphs: $r - 1$ strongly connected subgraphs with vertices $\{\chi_i, \chi_{\sigma(i)}\}_{i=1}^{r-1}$, and a strongly connected subgraph on the remaining vertices. $\square$

5.5. Three Jordan–Hölder factors: an example. In this section, we provide an example of an irreducible symplectic p -adic representation ρ for which Γ_ρ^{GSp} is an empty graph. This contrasts [Theorem 1.7](#). An important distinction of this example with the one provided in [Example 5.6](#) is that the example in [Example 5.6](#) is residually reducible over $\text{GL}_4(\mathbb{F}_p)$ and not over $\text{GSp}_4(\mathbb{F}_p)$. That is, the residual representation in [Example 5.6](#) does not stabilize an isotropic subspace.

PROPOSITION 5.14. *There exists an irreducible symplectic p -adic representation $\rho: G \rightarrow \text{GSp}_4(\mathbb{Z}_p)$ with three residual factors, satisfying (dist) and (rmf), such that Γ_ρ^{GSp} is the empty graph and Γ_ρ^{GL} is the complete graph.*

PROOF. Let $I_4 = \{1, 2, 3, 4\}$, and let $\mathcal{G} = (I_4, I_4^2, \eta)$ be the numbered graph defined by

- (i) $\eta_{i,i} = 0$ for all $i \in I_4$,
- (ii) $\eta_{i,j} = 1$ for all $1 \leq i \neq j \leq 3$,
- (iii) $\eta_{4,1} = \eta_{4,3} = \eta_{1,4} = \eta_{4,4} = 1$ and $\eta_{4,2} = \eta_{2,4} = 0$.

By definition, for all $i, j, k \in I_4$, $\eta_{ij} + \eta_{jk} \in \{0, 1, 2\}$. To show that η is a metric, the only case to check is when $\eta_{ij} + \eta_{jk} = 0$. In this case, it follows that $\eta_{ik} = 0$ by definition.

Let G be the group of elements $(a_{ij}) \in \text{GSp}_4(\mathbb{Z}_p)$ with $a_{ij} \in p^{\eta_{i,j}}\mathbb{Z}_p$. It is immediate that G is in fact the preimage in $\text{GSp}_4(\mathbb{Z}_p)$ under the reduction mod p map of the Levi subgroup $\text{GL}_1(\mathbb{F}_p) \times \text{GL}_1(\mathbb{F}_p) \times \text{GL}_2(\mathbb{F}_p)$ of the Klingen parabolic in $\text{GSp}_4(\mathbb{F}_p)$. Let $\rho: G \hookrightarrow \text{GSp}_4(\mathbb{Z}_p)$ be the inclusion. Then ρ is irreducible, residually multiplicity free with three residual factors, and satisfies (dist) by the definition of G . Let Σ be the apartment in $\mathcal{X}_4$ with respect to the chosen symplectic basis. By [\[BG06, Proposition 3.1.1\]](#), $\mathcal{S}_\rho^{\text{GL}}$ lies in the sub-apartment of Σ of dimension 2 in $\mathcal{X}_3$ and, by [\[BG06, Lemme 3.2.1\]](#), we have

$$\mathcal{S}_\rho^{\text{GL}} = \{[a_1, a_2, a_3, a_4] \in \Sigma \mid a_i - a_j \leq \eta_{i,j}\}.$$

By our choice of η , the simplicial complex $\mathcal{S}_\rho^{\text{GL}}$ is a triangulated hexagon in the sub-apartment of Σ . By [\[BG06, Théorème 3.3.3\]](#), Γ_ρ^{GL} is the complete graph on the three residual factors. On the other hand, we see that $S_\rho^{\text{GSp}} = \{[0, 0, 0, 0]\}$ by [Proposition 4.4](#). If Λ_0 is the primitive G -stable lattice such that $[\Lambda_0] = [0, 0, 0, 0]$, then $\bar{\Lambda}_0$ is semisimple and Γ_ρ^{GSp} has no edges. $\square$

References

- [AB08] Peter Abramenko and Kenneth S. Brown, *Buildings*, Graduate Texts in Mathematics, vol. 248, Springer, New York, 2008, Theory and applications. MR 2439729 [18](#), [30](#)
- [BC04] Joël Bellaïche and Gaëtan Chenevier, *Formes non tempérées pour $\text{U}(3)$ et conjectures de Bloch-Kato*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 4, 611–662. MR 2097894 [2](#)
- [BC09] Joël Bellaïche and Gaëtan Chenevier, *Families of Galois representations and Selmer groups*, Astérisque, no. 324, Société mathématique de France, 2009 (en). MR 2656025 [5](#), [11](#), [12](#), [13](#)
- [Bel03] Joël Bellaïche, *à propos d’un lemme de Ribet*, Rend. Sem. Mat. Univ. Padova **109** (2003), 45–62. MR 1997986 [2](#), [5](#), [18](#), [25](#), [28](#), [29](#), [30](#)
- [Bel09] Joël Bellaïche, *Ribet’s lemma, generalizations, and pseudocharacters*, Two lectures at the Clay Mathematical Institute Summer School, 2009. [2](#)
- [Bel19] Joël Bellaïche, *Image of pseudo-representations and coefficients of modular forms modulo p* , Adv. Math. **353** (2019), 647–721. MR 3989529 [16](#), [17](#)

- [BG06] Joël Bellaïche and Philippe Graftieaux, *Représentations sur un anneau de valuation discrète complet*, Math. Ann. **334** (2006), no. 3, 465–488. MR 2207872 [2](#), [4](#), [5](#), [11](#), [26](#), [34](#), [35](#), [36](#)
- [BK21] Tobias Berger and Krzysztof Klosin, *Irreducibility of limits of Galois representations of Saito-Kurokawa type*, Res. Number Theory **7** (2021), no. 3, Paper No. 41, 25. MR 4269673 [2](#)
- [Che14] Gaëtan Chenevier, *The p -adic analytic space of pseudocharacters of a profinite group and pseudorepresentations over arbitrary rings*, Automorphic forms and Galois representations. Vol. 1, London Math. Soc. Lecture Note Ser., vol. 414, Cambridge Univ. Press, Cambridge, 2014, pp. 221–285. MR 3444227 [5](#)
- [DK23] Samit Dasgupta and Mahesh Kakde, *On the Brumer-Stark conjecture*, Ann. of Math. (2) **197** (2023), no. 1, 289–388. MR 4513146 [5](#)
- [DKSW23a] Samit Dasgupta, Mahesh Kakde, Jesse Silliman, and Jiuya Wang, *The Brumer–Stark conjecture over $\mathbb{Z}$* , 2023. [5](#)
- [DKSW23b] ———, *The residually indistinguishable case of Ribet’s method for GL_2* , 2023. [5](#)
- [Gar97] Paul Garrett, *Buildings and classical groups*, Chapman & Hall, London, 1997. MR 1449872 [19](#)
- [GK96] Andrew V. Goldberg and Alexander V. Karzanov, *Path problems in skew-symmetric graphs*, Combinatorica **16** (1996), no. 3, 353–382. MR 1417346 [3](#)
- [LS10] T. Y. Lam and R. G. Swan, *Symplectic modules and von Neumann regular matrices over commutative rings*, Advances in ring theory, Trends Math., Birkhäuser/Springer Basel AG, Basel, 2010, pp. 213–227. MR 2664673 [6](#)
- [MQ26] Mohamed Moakher and Julian Quast, *Symplectic determinant laws and invariant theory*, Journal of the Institute of Mathematics of Jussieu **25** (2026), no. 5, 2853–2918. [5](#), [10](#), [11](#), [13](#)
- [MW84] B. Mazur and A. Wiles, *Class fields of abelian extensions of $\mathbb{Q}$* , Invent. Math. **76** (1984), no. 2, 179–330. MR 742853 [5](#)
- [Rib76] Kenneth A. Ribet, *A modular construction of unramified p -extensions of $\mathbb{Q}(\mu_p)$* , Invent. Math. **34** (1976), no. 3, 151–162. MR 419403 [1](#), [2](#)
- [Set08] Alison Setyadi, *The affine buildings of $SL(n)$ and $Sp(n)$: A combinatorial perspective*, ProQuest LLC, Ann Arbor, MI, 2008, Thesis (Ph.D.)—Dartmouth College. MR 2712416 [24](#)
- [She07] Thomas R. Shemanske, *The arithmetic and combinatorics of buildings for Sp_n* , Trans. Amer. Math. Soc. **359** (2007), no. 7, 3409–3423. MR 2299461 [24](#)
- [SU14] Christopher Skinner and Eric Urban, *The Iwasawa main conjectures for GL_2* , Invent. Math. **195** (2014), no. 1, 1–277. MR 3148103 [5](#)
- [Urb01] Eric Urban, *Selmer groups and the Eisenstein-Klingen ideal*, Duke Math. J. **106** (2001), no. 3, 485–525. MR 1813234 [3](#), [4](#), [5](#), [16](#), [18](#)
- [Wil90] A. Wiles, *The Iwasawa conjecture for totally real fields*, Ann. of Math. (2) **131** (1990), no. 3, 493–540. MR 1053488 [5](#)

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PA 19122, USA

Email address: aniruddha.sudarshan@temple.edu